



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/31, C07K 14/24, C12N 15/62, A61K 39/02	A1	(11) International Publication Number: WO 96/28551 (43) International Publication Date: 19 September 1996 (19.09.96)
(21) International Application Number: PCT/GB96/00571 (22) International Filing Date: 13 March 1996 (13.03.96) (30) Priority Data: 9505059.7 13 March 1995 (13.03.95) GB 9518946.0 15 September 1995 (15.09.95) GB 9524825.8 5 December 1995 (05.12.95) GB (71) Applicant (for all designated States except US): THE SECRETARY OF STATE FOR DEFENCE [GB/GB]; Defence Evaluation & Research Agency, DRA Farnborough, Hampshire GU14 6TD (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): TITBALL, Richard, William [GB/GB]; CBDE, Porton Down, Salisbury, Wiltshire SP4 0JQ (GB); WILLIAMSON, Ethel, Diane [GB/GB]; CBDE, Porton Down, Salisbury, Wiltshire SP4 0JQ (GB); LEARY, Sophie, Emma, Claire [GB/GB]; CBDE, Porton Down, Salisbury, Wiltshire SP4 0JQ (GB); OYSTON, Petra, Claire, Farquhar [GB/GB]; CBDE, Porton Down, Salisbury, Wiltshire SP4 0JQ (GB); BENNETT, Alice, Marie [GB/GB]; CBDE, Porton Down, Salisbury, Wiltshire SP4 0JQ (GB).	(74) Agent: BOWDERY, Anthony, Oliver, Defence Research Agency, Intellectual Property Dept., R69 Building, DRA Farnborough, Hampshire GU14 6TD (GB). (81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: VACCINES FOR PLAGUE		
(57) Abstract <p>A method of protecting a human or animal body from the effects of infection with <i>Y. pestis</i> is provided comprising administering to the body a vaccine including <i>Yersinia pestis</i> V antigen and <i>Yersinia pestis</i> F1 antigens or a protective epitopic part of each of these in a form other than whole <i>Y. Pestis</i> organisms. Preferably the antigens are administered in the form of a live vaccine or as recombinantly produced isolated and/or purified proteins. DNA encoding the whole or part of the F1 antigen and DNA encoding the whole or part of the V antigen may be used directly as a genetic vaccine.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MY	Malaysia
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Vaccines for plague

The present invention relates to novel vaccines for provision of protection against infection with the organism *Yersinia pestis* and to methods for administering these. Particularly provided are parenterally and orally active vaccines capable of offering protection against bubonic and pneumonic plague, particularly by induction of mucosal immunity in both humans and other animals.

Yersinia pestis is the highly virulent causative organism of plague in a wide range of animals, including man. Infection with such organisms results in a high rate of mortality. Studies have shown that the high virulence is due to a complex array of factors encoded by both the chromosome and three plasmids, including the Lcr genes, a fibrinolysin and a capsule.

Man is an occasional host in the natural cycle of the disease, and bubonic plague, characterised by the swelling of local lymph nodes, may occur following the bite of an infected flea. One of the complications of bubonic plague is secondary pneumonia, and in these cases the disease is readily transmitted between humans by airborne droplets. Plague is endemic in regions of North and South America, Africa, China and Asia (see Butler (1983) *Plague and Other Yersinia Infections*; Plenum Press, New York). Current outbreaks are believed to be part of the fourth world pandemic of the disease, with a clear need to protect individuals living or travelling in endemic areas, and laboratory workers handling the bacterium.

The current whole cell vaccines available for prevention of plague are highly heterogeneous, resulting in side effects which make them unsuitable for widespread use (eg Meyer et al (1974) *J. Infect Dis* 129 supp 13-18 and 85-120; Marshall et al (1974) *ibid* supp 19-25).

One current vaccine for plague is the Cutter vaccine which comprises formaldehyde killed plague bacilli and is administered to the body by intramuscular injection. However, parenteral immunisation, although effective in inducing systemic immunity, does not effectively induce mucosal immunity (McGhee et al, (1992) *Vaccine* 10, 75-88) and cases of pneumonic plague

have been reported in vaccinated individuals (Meyer (1970) Bull WHO 42 p653-666). So far no vaccine capable of producing a protective immune response at mucosal surfaces has been reported.

The live attenuated EV76 vaccine was tested extensively and used in the former Soviet Union from 1939, although its efficacy in evoking an immune response in man is questionable (Meyer et al (1974) J. Infect. Dis. 129 Supp: 13-18). It has been shown that the virulence of EV76 differs in several animal species, and non-human primates are particularly susceptible to a chronic infection with this strain. In the Western World the vaccine is considered to be unsuitable for mass vaccinations due to the extreme severity of the side effects and the possibility of the strain reverting to full virulence.

Two known Y. pestis antigens are the Y. pestis LcrV (V antigen), and the F1 antigen; both of which have now been found to be capable of evoking protective immune responses in animals. The present inventors have previously provided live orally active vaccine micro-organisms capable of expressing V antigen and F1 antigen respectively which each provide good protection against challenge with Y. pestis at up to 10^3 cfu. These vaccines are the subject of copending patent applications PCT/GB94/02818 and GB 9404577.0.

The present inventors have now surprisingly found that whereas only the unacceptably hazardous EV live vaccine had been shown to be capable of giving good protection against challenge with 10^8 cfu or more with Y. pestis GB strain, and V and F1 antigens alone only provide full protection against challenge with about 10^5 cfu, by administering a combined vaccine comprising V and F1 antigens they can at least match the protection afforded by EV76 without any of the hazards that have kept the EV vaccine from general use.

Still more advantageously, they have found that the vehicle for administration may be a simple mixture of the two protein components, rather than as a more complex attenuated whole organism. For long term protection and for the purposes of producing mucosal immunity, they have provided preferred forms of vaccine comprising the two components in the form of live attenuated vaccine such as the F1 and V expressing Aro A or C Salmonella

typhi referred to in the aforesaid depending applications, and in more preferred forms a single or double mutant expressing these antigens separately, or a fusion protein comprising both antigens.

Further provided are micro-organisms comprising both of F1 and V types of construct or plasmids of the applicants depending applications referred to above. These contain constructs that are capable of transforming a human or animal gut colonising micro-organism such that it is enabled to express proteins that are equivalent in sequence to F1 and V antigens respectively; these producing a protective immune response against Yersinia pestis in a human or animal body when the micro-organism is administered by oral or parenteral routes, and preferably allow the micro-organism to maintain its ability to colonise the human or animal gut.

A particularly preferred recombinant DNA, plasmid or human or animal gut colonising organism encodes for or expresses all or a protective epitopic part of the mature V protein of Yersinia pestis and all or a protective epitopic part of the mature F1 protein of Yersinia pestis. DNA encoding the whole or part of the F1 antigen and DNA encoding the whole or part of the V antigen could be used directly as a genetic vaccine.

Particularly preferred recombinant DNA encoding for V comprises a DNA sequence as described in SEQ ID No 1 or SEQ ID No 3, more preferably positioned in frame with a promoter such as *lacZ* or *nirB*, and preferably in a vector capable of expression and replication in a Salmonella. Particularly preferred recombinant DNA encoding for F1 comprises a DNA sequence as described in SEQ ID No 10. SEQ ID No 2 and SEQ ID No 4 show the amino acid sequences of two preferred V antigen proteins; SEQ ID No 2 being the sequence of the V-antigen itself, and SEQ ID No 4 being that of V-antigen with four extra vector defined N-terminal amino acids. SEQ ID No 11 is that of an F1 protein as encoded for by SEQ ID No 10.

The preferred DNA constructs used in micro-organisms of the invention allow production of micro-organisms that when orally administered induce local stimulation of the gut-associated

lymphoid tissue (GALT) and, by trafficking of lymphocytes through the common mucosal immune system provide a secondary stimulation of the bronchial associated lymphoid tissue (BALT). In this manner a secretory IgA response is achieved at the respiratory mucosal surface.

The micro-organisms provided by transformation using this DNA in vector or directly inserted format, are preferably attenuated, more preferably attenuated salmonella.

Attenuated micro-organisms such as S. typhimurium have been well characterised as carriers for various heterologous antigens (Curtiss, (1990); Cardenas and Clements, (1992)).

Attenuation may be effected in a number of ways, such as by use of the aro A and/or aro C mutation approach (see Hosieth et al (1981) Nature 291, 238-239, Dougan et al (1986) Parasite Immunol 9, 151-160; Chatfield et al (1989) Vaccine 7, 495-498); multiple mutations such as aro A and aro C mutants as described by Hone et al (1991) Vaccine 9, pp 810-816 may also be used. However, any suitably defective organism that is safe for intended use may be employed.

Many other such attenuated deletions and mutations will be known for these and other micro-organisms which will render them suitable for transformation with constructs of the present invention for the purposes of expressing vaccine proteins in the gut and/or gut colonisation in animals to be treated for Y. pestis. For human vaccination vectors containing the constructs of the present invention are placed in attenuated S. typhi and that transformed organism used as active agent for a live oral vaccine.

When DNA is used to transform the attenuated micro-organism by direct insertion into micro-organism DNA this may be by direct integration into a gene. Alternatively when incorporated in the form of a plasmid that expresses V or F1 protein or epitopic fragments thereof this may be such that only the V or F1 protein or fragment is expressed or that this is expressed as a fusion peptide with a further protein or peptide fragment, preferably including the other one of the antigenic F1/V components. Such further protein or peptide fragment might be such as

to promote export of mature proteins or peptide through the cell membrane or might be a further Y. pestis antigen.

The *lcr* gene was cloned from Y. pestis strain KIM by Price et al and its nucleotide sequence published in J Bacteriol (1989) 171, pp 5646-5653. In the examples below this information was used to design oligonucleotide primers which could amplify the gene from Y. pestis (strain GB) using the polymerase chain reaction (PCR). PCR primers were designed to be complementary to respective sequences flanking the 5' and 3' ends of the *lcrV* gene but also having 5' end tails containing a restriction enzyme recognition site to enable amplified *lcrV* gene to be cloned directionally into a plasmid vector (the 5' PCR primer containing an *EcoRI* site and the 3' primer containing a *SacI* site). These restriction enzyme sites are examples only and should not be seen as excluding other restriction enzymes.

In the examples below the constructs of the invention include a *lac* promoter, but other promoters such as the macrophage promoter (*nirB*) may be used.

The production of F1 has been described fully in Oyston et al (1995) Infect. Immun. Vol. 63 No 2 p563 - see page 564 under results: Cloning and Expression of *caf1*.

The dosage of the antigen components in a vaccine may vary dependent upon an individual animals immune characteristics, but for immunisation in the mouse animal model of the examples below it has been found that 10µg of each of V and F1 per dose were effective in providing full protection when administered in a standard primer and booster schedule.

The antigens may be incorporated into a conventional pharmaceutically acceptable carrier, no particular limitation being imposed here. Conveniently the antigens have been incorporated into an oil in water emulsion. Adjuvants may be included in the vaccine composition, and particularly Freund's Incomplete adjuvant IFA has been found to be effective when treating the mouse model.

The carrier may be one suited to parenteral administration, particularly intraperitoneal administration but optionally oral, in the case of micro-organism based vaccines, or administration in the form of droplets or capsules, such as liposomes or microcapsules as would be effective in delivering the composition to the airways of an individual for the purpose of evoking mucosal immune response. The carrier may also comprise a slow-depot release system e.g. Alhydrogel.

Another method of encapsulation includes the use of polymeric structures in particular linear block co-polymers. Biodegradable polymers for example poly-lactic acid with or without glycolic acid or block co-polymer may be used; these may contain the following repeat unit: $(POP-POE)_n$ where POP is polyoxypropylene and POE is polyoxyethylene. Block co-polymers which contain $(POP-POE)_n$ are of particular use.

The method, constructs, micro-organisms and vaccines of the invention will now be exemplified by way of illustration only by reference to the following Sequence listing, Figure and Examples. Still further embodiments will be evident to those skilled in the art in the light of these.

Figure 1 illustrates in bar-chart form the survival rates of a number of groups against a challenge of *Y.pestis*.

Figure 2 illustrates in graphical form, IgG priming responses to intramuscular BSA immunisation in Balb/c mice.

SEQUENCE LISTING

SEQ ID No 1: Shows the nucleotide and derived amino acid sequence of a V-encoding DNA with 6 bases of vector pMAL-p2 or pMAL-c2 into which it is cloned at the 5' end using the EcoRI site in sequence GAATTC (derived from the 5' end PCR primer) and at the 3' end at the SalI site in sequence GTCGAC (derived from the 3' end PCR primer). The base at position 1006 has been altered by PCR mutagenesis to a T to create a second in frame stop codon. The start of the amino acid sequence is C-terminal to a factor Xa cleavage site.

SEQ ID No 2: Shows the amino acid sequence of the peptide expressed by the insert DNA of the invention, with an additional four amino acids encoded for by the vector (I E + F S) at the N-terminal end.

SEQ ID No 3: Shows the nucleotide and derived amino acid sequence of a second V-encoding DNA of the invention with 10 bases of a vector pGEX-5X-2 into which it is cloned shown at the 5' end using the EcoRI site in sequence GAATTC (GA derived from the 5' end PCR primer) and the SalI site in sequence GTCGAC (GTCGAC derived from the 3' end PCR primer). The base at position 1006 has been altered by PCR mutagenesis to create a second in frame stop codon; the base at position 16 has been altered to a C from an A to create the EcoRI site. The start of the amino acid sequence is C-terminal to a factor Xa cleavage site.

SEQ ID No 4: Shows the amino acid sequence of the peptide expressed by the DNA of SEQ ID No 3, with four amino acids encoded by the vector (G, I, P and G) at the N-terminal end.

SEQ ID No 5: Shows the nucleotide sequence of a gene 5' end primer oligonucleotide used to generate V-encoding DNA used in SEQ ID No 1.

SEQ ID No 6: Shows the nucleotide sequence of a gene 3' end primer oligonucleotide used to generate V-encoding DNA used in the Examples.

SEQ ID No 7: Shows the nucleotide sequence of a PCR primer oligonucleotide corresponding to the first 21 bases encoding for mature caf1 with an additional 5' region encoding for a SacI site.

SEQ ID No 8: Shows the nucleotide sequence of a PCR primer oligonucleotide corresponding to the sequence of caf1 which encodes a 'stem loop' downstream of the termination codon with an added 5' region encoding SacI and AclI sites.

SEQ ID No 9: Shows the nucleotide sequence of a PCR primer oligonucleotide corresponding to an internal end region of the caf1 gene starting 107 bases downstream from the end of the first oligonucleotide.

SEQ ID No 10: Shows the nucleotide sequence of the pFGAL2a construct showing the fusion of the first few bases of the β -galactosidase sequence in the vector with *caf1* minus its signal sequence and having a 5' tail including a *Sac* I restriction site; the sequence is shown up to the *caf1* AACC 3' end with some vector bases.

SEQ ID No 11: Shows the amino acid sequence of the protein encoded by pFGAL2a. This sequence may be preceded by Met, Thr, Met, Ile, Thr, Asn.

SEQ ID No 12: is that of primer F1OU2 used to amplify the F1 operon.

SEQ ID No 13: is that of primer M4D used to amplify the F1 operon.

SEQ ID No 14: is that of primer M3U used to amplify the F1 operon.

SEQ ID No 15: is that of primer F1OD2 used to amplify the F1 operon.

SEQ ID No 16: is the nucleotide and derived amino acid sequence of a DNA fragment encoding an F1-V fusion protein. There is a *Sac*I cloning site at the 5' end and a *Hind* III cloning site at the 3' end. Bases 452-472 is a sequence contained in the cloned insert, but derived from PCR primers (not found in *Y. pestis* DNA).

SEQ ID No 17: is the amino acid sequence of SEQ ID No 16.

SEQ ID No 18: is that of primer 5'FAB2 used to amplify the F1 operon including signal sequence.

SEQ ID No 19: is that of primer 3'FBAM used to amplify the F1 operon including signal sequence.

SEQ ID No 20: is the nucleotide and amino acid sequence for F1 antigen as defined by PCR primers detailed in exemplified SEQ ID No 18 and 19 including signal sequence.

SEQ ID No 21: is the amino acid sequence of SEQ ID No 20.

SEQ ID No 22: is the nucleotide and amino acid sequence of F1/V fusion protein including a 3 amino acid linker region. The T at position 1522 was modified from G to create a second in frame stop codon.

SEQ ID No 23: is the amino acid sequence of SEQ ID No 22. The linker region referred to in SEQ ID No 22 is at amino acid position 171-176 (bases 523-540 in SEQ. ID No 22).

SEQ ID No 24: shows the nucleotide sequence of a gene 5' end primer oligonucleotide used to generate V-encoding DNA used in SEQ ID No 3.

EXAMPLES

Cloning of SEQ ID No 3

Materials and Methods: Materials for the preparation of growth media were obtained from Oxoid Ltd. or Difco Laboratories. All enzymes used in the manipulation of DNA were obtained from Boehringer Mannheim UK Ltd. and used according to the manufacturers instructions. All other chemicals and biochemicals were obtained from Sigma chemical Co unless otherwise indicated. Monospecific rabbit polyclonal anti-V and mouse anti-GST sera were prepared by Dr R Brubaker (Department of Microbiology, Michigan State University) and Dr E D Williamson (Chemical and Biological Defence Establishment), respectively.

Bacterial strains and cultivation: *Yersinia pestis* GB was cultured aerobically at 28°C in a liquid medium (pH 6.8) containing 15g proteose peptone, 2.5g liver digest, 5g yeast extract, 5g NaCl per litre supplemented with 80ml of 0.25% haemin dissolved in 0.1M NaOH (Blood Base Broth). *Escherichia coli* JM109 was cultured and stored as described by Sambrook et al. Molecular Cloning. A Laboratory Manual.

Production of recombinant V and F1 proteins: Manipulation of DNA. Chromosomal DNA was isolated from *Y. pestis* by the method of Marmur.

Production of recombinant V-antigen: The gene encoding V-antigen (1crV) was amplified from *Y. pestis* DNA using the polymerase chain reaction (PCR) with 125pmol of primers homologous to sequences from the 5' and 3' ends of the gene (see Price et al (1989). J. Bacteriol 171 p5646-5653).

The sequences of the 5' primer (V/5'E: GATCGAATTCGAGCCTACGAACAA) and the 3' primer (GGATCGTCGACTTACATAATTACCTCGTGTCA) also included 5' regions encoding the restriction sites EcoRI and SalI, respectively. In addition, two nucleotides were altered from the published sequence of 1crV (Price et al, 1989), so that the EcoRI site was completed and the amplified gene encoded an extra termination codon (TAA). The PCR primers were

prepared with a DNA synthesiser (392 Applied Biosystems) Applied Biosystems. A DNA fragment was obtained after 30 cycles of amplification (95°C, 20secs, 45°C, 20secs, 72°C, 30secs; Perkin 9600 GensAmp PCR System). The fragment was purified, digested with EcoRI and SalI, ligated with suitably digested plasmid pGEX-5X-2 and transformed into *E. coli* JM109 by electroporation (see Sambrook et al 1989). A colony containing the recombinant plasmid (pVG100) was identified by PCR using 30-mer primers (5' nucleotides located at positions 54 and 794; see Price et al 1989) which amplified an internal segment of the *lcrV* gene.

Expression of rV in *E. coli*. Cultures of *E. coli* JM109/pVG100 were grown in LB containing 100µgml⁻¹ ampicillin at 37°C until the absorbance (600nm) was 0.3. Isopropyl-β-D-thiogalactopyranoside (IPTG) was then added to the culture to a final concentration of 1mM and growth was continued for a further 5 hours. Whole cell lysates of the bacteria were prepared as described in Sambrook et al and expression of the GST/V fusion protein was examined by staining 10% SDS-polyacrylamide gels (Mini-Protean II, BioRad) with Coomassie Brilliant Blue R250 and by Western Blotting (see Sambrook et al). Western Blots were probed with rabbit anti-V serum diluted 1/4000 or mouse anti-GST serum in dilution at 1/1000 and protein bands were visualised with a colloidal gold labelled secondary antibody (Auroprobe BLplus, Cambio).

Quantification of GST/V expression in vitro. Cultures of *E. coli* JM109 containing pVG100 or pGEX-5X-2 were grown as described above. One ml aliquots were removed from the cultures in logarithmic and stationary phases and the number of viable cells determined by inoculating onto L-agar containing ampicillin. The cells were harvested from a second 1 ml aliquot by centrifugation and resuspended in 1 ml of phosphate buffered saline (PBS). The cell suspension was frozen at -20°C for 1 hour, thawed and then sonicated on ice at 10°C lower for 3 x 30 secs (model XL2015 sonicator, 3.2mm Microtip probe; Heat Systems Inc.). The sonicates and a standard solution of rV (5µgml⁻¹) were serially diluted in PBS in a microtitre plate and allowed to bind overnight at 4°C. The quantity of GST/V fusion protein in each sonicate was determined in a standard ELISA using rabbit anti-V serum as the primary antibody. Antibodies were incubated in 1% skimmed milk powder in PBS.

Purification of rV. *E. coli* JM109/pVG100 was grown in 5 x 100 ml volumes of LB as described above. The cells were harvested by centrifugation and resuspended in 3 ml phosphate buffered saline (PBS). After the addition of 80ul lysozyme (10 mgml⁻¹), the cell suspension was incubated for 10 min at 22°C. Triton X-100 was added to a final concentration of 1% and the cells were frozen (-20°C), thawed and sonicated on ice at 70% power for 3 x 30 s (model XL2015 sonicator). The lysed cells were centrifuged, and the supernatant was made up to 30ml with PBS and mixed with 5ml of Glutathione Sepharose 4B (Pharmacia Biotech) which had been washed three times with PBS + 0.1% Triton X-100. The mixture was stirred for 18 hours at 4°C, centrifuged and washed twice in 100 ml PBS, and then packed into a chromatography column (Poly-Prep; Bio-rad) as a 50% slurry. The GST/V fusion protein was eluted with 10 ml of 50 mM Tris pH 8.0 containing 5 mM reduced glutathione (Pharmacia Biotech). After dialysis against PBS, the fusion protein was cleaved with factor Xa (Boehringer Mannheim UK Ltd) for 18 hours at 22°C, according to the manufacturer's instructions. Cleaved GST and excess uncleaved GST/V were removed from the solution by affinity adsorption, as described above, to leave purified recombinant V (rV).

Immunisation with rV. Six week old female Balb/c mice, raised under specific pathogen-free conditions (Charles. River Laboratories, Margate, Kent, UK), were used in this study. A group of 16 mice received a 0.2ml primary immunising dose intraperitoneally (i.p.) of 10.13µg of rV antigen, presented in a 1:1 water-in-oil emulsion with Incomplete Freund's Adjuvant (IFA). On days 14 and 34, each animal received booster immunising doses, prepared as above. On day 64, 6 animals were sacrificed and their tissues were removed for immunological analyses, as described below. The remaining animals were challenged with *Y. pestis*. An untreated control group of 16 age-matched mice were divided similarly into groups for tissue sampling and challenge. In a subsequent experiment to determine the degree of protection against higher challenge doses of *Y. pestis* groups of 5 or 6 rV-immunised and control mice were used.

Measurement of serum antibody titre. Blood was sampled by cardiac puncture from mice anaesthetised i.p. with a 0.1ml cocktail containing 6mg of Domitor (Norden Laboratories) and

27µg of Ketaiar (Parks-Davies). The samples were pooled and the serum was separated. The serum antibody titre was measured by a modified ELISA (Williamson and Tiball, (1993) Vaccine 11: 1253-1258). Briefly, rV (5 µgml⁻¹ in PBS) was coated onto a microtitre plate and the test sera were serially diluted in duplicate on the plate. Bound antibody was detected using peroxidase labelled conjugates of anti-mouse polyvalent Ig. The titre of specific antibody was estimated as the maximum dilution of serum giving an absorbance reading greater than 0.1 units, after subtraction of the absorbance due to non-specific binding detected in the control sera.

Isolation of purified T cells from the spleen. A crude suspension of mixed spleen cells was prepared by gently grinding the spleen on a fine wire mesh. The cells were flushed from the splenic capsule and connective tissue with 2ml of tissue culture medium (DMEM with 20mM L-glutamine, 10⁵U l⁻¹ of penicillin and 100mg l⁻¹ of streptomycin). A population of mixed lymphocytes was separated from the spleen cell suspension by density gradient centrifugation of Ficoll-Hypaque (Lymphocyte Separation Medium, ICN Flow). A mixed acridine orange (0.0003% w/v) and ethidium bromide (0.001% w/v) stain was used to determine the percentage of viable cells in the preparation.

The mixed lymphocytes were incubated with sheep anti-mouse IgG-coated Dynabeads (M450, Dynal UK) at a ratio of 1:3 for 30 minutes at 4°C. The Dynabead linked B cells were removed by magnetic separation and the remaining T cells were resuspended in DMEM, supplemented with antibiotic and 10% v/v foetal calf serum (FCS) at the desired cell density for seeding to microtitre plates.

In vitro proliferation of crude spleen cells and purified T cells against rV. Doubling dilutions of rV or Concanavalin A (positive control) in DMEM (range 0-50µgml⁻¹) were made in the wells of a microtitre plate, such that 0.1ml remained in each well. Negative controls consisted of 0.1ml of DMEM alone. An equal volume of the crude spleen cell or purified T cell suspension was seeded into each well at a minimum density of 5x10⁴ cells and incubated for 72 hours at 37°C (5% CO₂). One µCi of ³H thymidine ([methyl-³H]thymidine S.A. 74 GBqmmol⁻¹; Amersham) in 30µl of DMEM supplemented with 10% FCS was aliquoted into each well and

incubation was continued for 24 h. The well contents were harvested onto glass fibre filters using a cell harvester (Titertek) and discs representing each well were punched from the filter mats into 1.5ml of scintillation fluid (Cytoscint, ICN Biomedicals Inc.) to measure the incorporation of ^3H thymidine into cells. The cell stimulation index was calculated from 4 replicates as mean cpm (stimulated)/mean cpm (negative control)

Production of recombinant F1 antigen. Cloning of *caf1*: DNA was isolated from *Y. pestis* by the method of Marmur et al (1961) J. Mol. Biol. 3: pp 208-218. A DNA fragment encoding the open reading frame of *caf1* minus its signal sequence was amplified from this using the polymerase chain reaction (PCR). Oligonucleotides were prepared with a Beckman 200A DNA synthesiser for use in the PCR.

pFGAL2a construct. Oligonucleotide GATCGAGCTCGGCAGATTAACTGCAAGCACC (SEQ ID No 7) was synthesised corresponding to the first 21 bases of the *caf1* gene immediately following the nucleotides encoding the signal sequence with an additional 5' region encoding a *SacI* site and the complimentary oligonucleotide CAGGTGCGAGCTCGTCGACGGTTAGGCTCAAAGTAG (SEQ ID No 8) corresponding to the sequence which encodes a putative 'stem loop' structure downstream of the *caf1* termination codon with an added 5' region encoding *SacI* and *AccI* sites. A DNA fragment was obtained after 35 cycles of amplification (95°C, 15 secs; 50°C, 15 secs; 72°C, 30 secs using a Perkin Elmer 9600 GeneAmp PCR system). The fragment was purified, digested with *SacI* and *AccI*, ligated into a similarly digested pUC18 plasmid and transformed into *E. coli* JM109 by electroporation. Electroporation was carried out using a Biorad Gene Pulser with 0.2 cm cuvettes at 1.25kV, 25µF, 800 Ohms with a time constant of 20.

A pFGAL2a colony containing the cloned *caf1* gene was identified by PCR using an oligonucleotide TGGTACGCTTACTCTTGGCGGCTAT (SEQ ID No 9) corresponding to an internal region of the gene 128 to 153 nucleotides from the site identified as the signal sequence cleavage site (see Gaiyov et al (1990)) and the SEQ ID No 2. An F1 expressing *E. coli* culture containing the pFGAL2a was grown at 37°C with shaking in Luria Broth with 1mM

isopropyl- β -D-thiogalactopyranoside (IPTG) for 18 hours. Whole cell lysates and periplasmic and cytoplasmic fractions of the bacteria were prepared as described by Sambrook et al (1989).

SDS-PAGE and Western blotting: SDS-polyacrylamide gel electrophoresis (PAGE) and Western blotting were performed as described by Hunter et al (1993) *Infect. Immun.* 61, 3958-3965. Blots were probed with polyclonal antisera raised in sheep (B283) against killed *Y. pestis* (EV76 strain grown at 37°C) and bound antibody was detected with a horseradish peroxidase-labelled donkey anti-sheep IgG (Sigma).

EXAMPLES 1 AND 2 OF COMBINATION V/F1 VACCINE. COMPARATIVE EXAMPLES.
EFFECT OF VACCINES ON MORTALITY OF MICE ON CHALLENGE WITH γ PESTIS
(GB) STRAIN

Animals: Barrier bred female 6-week old Balb/c mice free of mouse pathogens were obtained from Charles-Reiver Laboratories, Margate, Kent, UK and were used throughout these Examples.

Immunisation: Mice were divided into groups of 5 or 6 and immunised as follows.

Comparative vaccine EV75: A total of 50 mice received a single subcutaneous (s.c.) priming dose of live EV75 vaccine on day 0 of the schedule delivered in a total volume of 100 μ l.

Comparative vaccine Cutter USP vaccine: A further group of 50 mice were primed intramuscularly (i.m.) with 100 μ l of Cutter vaccine; this representing about 0.2 of the human dose and comprised approximately 2×10^8 formaldehyde killed plague bacilli. This dose was administered to each animal again on day 16 of the schedule to effect booster immunisation.

F1 and V vaccines: Groups of 62 mice received a primary immunising dose intra-peritoneally (i.p.) of either recombinant V-antigen or recombinant F1-antigen, presented in a 1:1 water-in-oil emulsion with incomplete Freund's adjuvant (IFA; Sigma). Animals were primed with a 10 μ g dose of the respective antigen in a total volume of 0.1ml emulsion. Animals were boosted with the respective antigen as appropriate on days 14 (V and F1 groups) and 28 (all F1 group and a sub-group of 12 of the V group).

A further group of 12 mice were primed and boosted on days 0, 14 and 28 with a combination of 10 μ g F1 and 10 μ g V jointly incorporated into the aqueous phase of a 1:1 water-in-oil emulsion with IFA (final volume 0.1ml per mouse).

On day 50 of the immunisation schedule, 6 animals were selected at random from each of the treatment groups for subsequent analysis of spleen cell responses. The remaining animals in each group were challenged with Y. pestis. An untreated control group of age-matched mice was similarly split.

Multiple LD challenge to determine limits of protection: Mice from each of the immunised groups and untreated controls were divided into groups of 5 or 6 for challenge by the s.c. route with Y. pestis GB strain in the dose range 20 to 2×10^8 viable organisms. Challenged mice were closely observed over a 14-day period for the development of symptoms and where appropriate time to death was carefully recorded.

Animals which succumbed to the challenge were autopsied and blood smears, livers and spleens removed for bacteriological analysis.

The results of these challenges on control and test animals is given in Table 1 below, two sets of results being given corresponding two experimental runs with respective controls. From these results it can clearly be seen that while the V antigen is more effective than Cutter, it is inferior to EV76. However, when combined with the less effective F1 the combination is as effective as EV76 without side effects.

TABLE 1

VACCINE	GB CHALLENGE (cfu)	SURVIVORS	MEAN TTD (hr±sem)	SKIN LESIONS	SICK
<u>Set 1 results</u>					
EV	2×10^9	5/5		+	-
EV	2×10^8	4/5		+	-
EV	2×10^7	6/6		+	-
EV	2×10^6	6/6		+	-
EV	2×10^5	6/6		+	-
Cutter	2×10^9	0/5	119±zero	+	-
Cutter	2×10^8	0/5	171±11.63	+	+
Cutter	2×10^7	1/5	160±zero	+	+
Cutter	2×10^6	3/5	120±8.49	+	+
V	2×10^8	0/5	102.4±21.4	+	+
V	2×10^6	4/5	156±zero	-	-
V	2×10^7	5/5		-	-
V	2×10^6	4/5	64±zero	-	-
V	2×10^5	4/5	112±zero	-	-
Control	2×10^9	0/5	64±zero	-	+
Control	2×10^4	0/5	121.6±16.08	-	+

VACCINE	GB CHALLENGE (cfu)	SURVIVORS	MEAN TTD (hr±sem)	SKIN LESIONS	SICK
<u>Set 2 results</u>					
F1	2×10^6	0/5	98.6±8.41	-	+
F1	2×10^7	3/5	124±8.49	-	+
F1	2×10^5	4/5	136±zero	-	+/-
F1	2×10^3	5/5		-	-
F1	20	5/5		-	-
V	2×10^8	0/5	102.4±8.59	-	+
V	2×10^5	5/5		-	-
F1+V*	2×10^6	5/5		-	-
F1+V**	2×10^5	5/5		-	-
Control	2×10^8	0/5	64±zero	-	+

* = Example 1

** = Example 2

EXAMPLE 3. Production of Attenuated Salmonella for use in oral vaccine.Expression of recombinant V-antigen from *S. typhimurium* and *typhi*.

Amplified *lcrV* gene was cloned into three different plasmid vectors:

pMAL-p2: a vector designed to express the cloned gene as a fusion product with a maltose binding protein (MBP). The C-terminus of the MBP is fused to the N-terminus of the V-antigen. The fusion protein so produced on expression is exported to the periplasm. Vector including the V-antigen DNA sequence was designated pVMP100.

pMAL-c2: a vector similar to pMAL-p2 except that MBP-V antigen fusion protein is expressed cytoplasmically. The recombinant plasmid was designated pVMC100.

pGEX-5X-2: a vector designed to express the cloned gene as a fusion protein with glutathione-S-transferase (GST). The C-terminus of GST is fused to the N-terminus of V antigen and the fusion protein is expressed cytoplasmically. The recombinant plasmid was designated pVG100.

All the vectors contain the P_{lac} promoter and the $lacI^Q$ gene; the latter encoding the lac repressor which turns off transcription from P_{lac} in *Escherichia coli* until IPTG is added. The plasmids contain the origin of replication from pBR322 and as a result replicate to a low copy number in the bacterial cell.

Each of the recombinant plasmids was electroporated into *Salmonella typhimurium* strain SL3261, an attenuated strain that has been used extensively as a live vaccine vector for the expression of foreign antigens. It contains a specific deletion mutation in the *aroA* gene which makes the mutant dependent upon certain aromatics for growth (see Hosie et al). For producing microorganism suitable for human vaccination use electroporation is into attenuated *Salmonella typhi*.

The recombinant plasmids all expressed V antigen as shown by Western blotting of *S. typhimurium* cultures and probing with a monospecific anti-V antigen polyclonal antiserum supplied by R Brubaker, Dept Microbiology, Michigan State University, East Lansing, MI 48824-1101, USA. Recombinant *S. typhimurium* were inoculated intravenously into mice at 5×10^7 cfu/dose and shown to colonise the liver and spleen at high levels: between 8×10^6 and 5×10^8 cfu per organ were recovered. The majority of the bacterial cells recovered were also ampicillin resistant indicating retention of recombinant plasmids.

Expression of F1 in *S. typhimurium*: The pFGAL2a plasmid was isolated using general techniques described in Sambrook et al (1989) Molecular Cloning; a Laboratory Manual, 2nd Edition. Cold Spring Harbour Laboratory, New York. Purified plasmid was electroporated into *S. typhimurium* LB5010 (restriction⁻, modification⁻) and methylated pFGAL2a was subsequently isolated from the LB5010 for electroporation into *S. typhimurium* SL3261 (aro⁻ A⁻). Periplasmic and cytoplasmic fractions were prepared for SDS-PAGE and Western blotting as described above.

Stability of constructs: Five female Balb/c mice were inoculated intravenously with either 5×10^5 or 5×10^7 cfu *S. typhimurium* containing pFGAL2a in 200 μ L phosphate buffered saline. Control mice were inoculated similarly with *S. typhimurium* containing pUC18 with no insert. After 7 days the mice were killed by cervical dislocation and their livers and spleens removed. The organs were homogenised in 10ml phosphate buffered saline using a stomacher on maximum setting for 2 minutes and the homogenate was serially diluted in phosphate buffered saline and placed onto L agar or L agar containing 55 μ g ml⁻¹ ampicillin.

F1 operon construct: Attempts to PCR replicate the entire F1 operon as one piece were unsuccessful, so a strategy was developed whereby it was amplified using PCR to produce two discrete fragments using primer pairs (A) of SEQ ID No 12 and 13 and (b) of SEQ No 14 and 15 respectively to produce fragments of 3.36kb and 1.89kb from *Y. pestis* MP6 template DNA. Marmur extract of DNA was used without CsCl₂ purification. The PCR cycle conditions used were 96°C for 30 seconds, 57°C for 30 seconds and 72°C for 1 minute; total of 30 cycles.

These two fragments were digested using *Nhe*I and joined together.

This fused fragment, encoding the full length operon (5.25kb), was digested with *Eco*R1 and *Sal*I and then cloned into a number of vectors. When this fragment was cloned into pBR322 and expressed in *E. coli*, *S. typhimurium* LB5010 or SL3261 instability of the recombinant plasmid was noted. To circumvent this problem the operon was cloned into plasmid pLG339, a low copy number plasmid km^R . The entire F1 operon was also inserted into *ArgC* gene on the chromosome of gene on the chromosome of *S. typhimurium* using vector pBRD1084.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: SECRETARY OF STATE FOR DEFENCE, ...

(ii) TITLE OF INVENTION: VACCINES

(iii) NUMBER OF SEQUENCES: 24

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: SECRETARY OF STATE FOR DEFENCE

(B) STREET: WHITEHALL

(C) CITY: LONDON

(D) STATE: LONDON

(E) COUNTRY: UNITED KINGDOM (GB)

(F) ZIP: SW1A 2HB

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

25

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1014 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:

- (A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..987

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATT TCA GAA TTC ATT AGA GCC TAC GAA CAA AAC CCA CAA CAT TTT ATT	48
Ile Ser Glu Phe Ile Arg Ala Tyr Glu Gln Asn Pro Gln His Phe Ile	
1 5 10 15	
GAG GAT CTA GAA AAA GTT AGG GTG GAA CAA TTT ACT GGT CAT GGT TCT	96
Glu Asp Leu Glu Lys Val Arg Val Glu Gln Leu Thr Gly His Gly Ser	
20 25 30	

TCA GTT TTA GAA GAA TTG GTT CAG TTA GTC AAA GAT AAA AAT ATA GAT 144
 Ser Val Leu Glu Glu Leu Val Gln Leu Val Lys Asp Lys Asn Ile Asp

38

40

45

ATT TCC ATT AAA TAT GAT CCC AGA AAA GAT TCG GAG GTT TTT GCC AAT 192
 Ile Ser Ile Lys Tyr Asp Pro Arg Lys Asp Ser Glu Val Phe Ala Asn

50

55

60

AGA GTA ATT ACT GAT GAT ATC GAA TTG CTC AAG AAA ATC CTA GCT TAT 240
 Arg Val Ile Thr Asp Asp Ile Glu Leu Leu Lys Lys Ile Leu Ala Tyr

65

70

75

80

TTT CTA CCC GAG GAT GCC ATT CTT AAA GGC GGT CAT TAT GAC AAC CAA 288
 Phe Leu Pro Glu Asp Ala Ile Leu Lys Gly Gly His Tyr Asp Asn Gln

85

90

95

CTG CAA AAT GGC ATC AAG CGA GTA AAA GAG TTC CTT GAA TCA TCG CCG 336
 Leu Gln Asn Gly Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro

100

105

110

AAT ACA CAA TGG GAA TTG CGG GCG TTC ATG GCA GTA ATG CAT TTC TCT 384
 Asn Thr Gln Trp Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser

115

120

125

TTA ACC GCC GAT CGT ATC GAT GAT GAT ATT TTG AAA GTG ATT GTT GAT 432
 Leu Thr Ala Asp Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp

130

135

140

TCA ATG AAT CAT CAT GGT GAT GCC CGT AGC AAG TTG CGT GAA GAA TTA 480
 Ser Met Asn His His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu
 145 150 155 160

GCT GAG CTT ACC GCC GAA TTA AAG ATT TAT TCA GTT ATT CAA GCC GAA 528
 Ala Glu Leu Thr Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu
 165 170 175

ATT AAT AAG CAT CTG TCT AGT AGT GGC ACC ATA AAT ATC CAT GAT AAA 576
 Ile Asn Lys His Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys
 180 185 190

TCC ATT AAT CTC ATG GAT AAA AAT TTA TAT GGT TAT ACA GAT GAA GAG 624
 Ser Ile Asn Leu Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu
 195 200 205

ATT TTT AAA GCC AGC GCA GAG TAC AAA ATT CTC GAG AAA ATG CCT CAA 672
 Ile Phe Lys Ala Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln
 210 215 220

ACC ACC ATT CAG GTG GAT GGG AGC GAG AAA AAA ATA GTC TCG ATA AAG 720
 Thr Thr Ile Gln Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys
 225 230 235 240

GAC TTT CTT GGA AGT GAG AAT AAA AGA ACC GCG GCG TTG GGT AAT CTG 768
 Asp Phe Leu Gly Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu
 245 250 255

28

AAA AAC TCA TAG TCT TAT AAT AAA GAT AAT AAT GAA TTA TCT CAC TTT 816

Lys Asn Ser Tyr Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe

260

265

270

ACC ACC ACC TCG TCG GAT AAG TCC AGG CCG CTC AAC GAC TTG GTT AGC 864

Ala Thr Thr Cys Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser

275

280

285

CAA AAA ACA ACT CAG CTG TCT GAT ATT ACA TCA CGT TTT AAT TCA GCT 912

Gln Lys Thr Thr Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala

290

295

300

ATT GAA GCA CTG AAC CGT TTC ATT CAG AAA TAT GAT TCA GTG ATG CAA 960

Ile Glu Ala Leu Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln

305

310

315

320

CGT CTG CTA GAT GAC ACG TCT GGT AAA TGACACGAGG TAATTATGTA 1007

Arg Leu Leu Asp Asp Thr Ser Gly Lys

325

AGTCGAC

1014

29

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 329 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ile	Ser	Glu	Phe	Ile	Arg	Ala	Tyr	Glu	Gln	Asn	Pro	Gln	His	Phe	Ile
1				5				10						15	
Glu	Asp	Leu	Glu	Lys	Val	Arg	Val	Glu	Gln	Leu	Thr	Gly	His	Gly	Ser
			20					25						30	
Ser	Val	Leu	Glu	Glu	Leu	Val	Gln	Leu	Val	Lys	Asp	Lys	Asn	Ile	Asp
			35				40							45	
Ile	Ser	Ile	Lys	Tyr	Asp	Pro	Arg	Lys	Asp	Ser	Glu	Val	Phe	Ala	Asn
			50				55							60	
Arg	Val	Ile	Thr	Asp	Asp	Ile	Glu	Leu	Leu	Lys	Lys	Ile	Leu	Ala	Tyr
			65			70				75				80	
Phe	Leu	Pro	Glu	Asp	Ala	Ile	Leu	Lys	Gly	Gly	His	Tyr	Asp	Asn	Gln
						85				90				95	

30

Leu Gln Asn Gly Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro
100 105 110

Asn Thr Gln Trp Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser
115 120 125

Leu Thr Ala Asp Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp
130 135 140

Ser Met Asn His His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu
145 150 155 160

Ala Glu Leu Thr Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu
165 170 175

Ile Asn Lys His Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys
180 185 190

Ser Ile Asn Leu Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu
195 200 205

Ile Phe Lys Ala Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln
210 215 220

Thr Thr Ile Gln Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys
225 230 235 240

Asp Phe Leu Gly Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu
245 250 255

31

Lys Asn Ser Tyr Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe
260 265 270

Ala Thr Thr Cys Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser
275 280 285

Gln Lys Thr Thr Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala
290 295 300

Ile Glu Ala Leu Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln
305 310 315 320

Arg Leu Leu Asp Asp Thr Ser Gly Lys
325

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1014 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..987

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```
GGG ATC CCC GGA ATT CGA GCC TAC GAA CAA AAC CCA CAA CAT TTT ATT   48
Gly Ile Pro Gly Ile Arg Ala Tyr Glu Gln Asn Pro Gln His Phe Ile
1             5             10             15
```

33

GAG GAT CTA GAA AAA GTT AGG GTC GAA CAA CTT ACT GGT CAT GGT TCT 95
 Glu Asp Leu Glu Lys Val Arg Val Glu Gln Leu Thr Gly His Gly Ser
 20 25 30

TCA GTT TTA GAA GAA TTG GTT CAG TTA GTC AAA GAT AAA AAT ATA GAT 144
 Ser Val Leu Glu Glu Leu Val Gln Leu Val Lys Asp Lys Asn Ile Asp
 35 40 45

ATT TCC ATT AAA TAT GAT CCC AGA AAA GAT TCG GAG GTT TTT GCC AAT 192
 Ile Ser Ile Lys Tyr Asp Pro Arg Lys Asp Ser Glu Val Phe Ala Asn
 50 55 60

AGA GTA ATT ACT GAT GAT ATC GAA TTG CTC AAG AAA ATC CTA GCT TAT 240
 Arg Val Ile Thr Asp Asp Ile Glu Leu Leu Lys Lys Ile Leu Ala Tyr
 65 70 75 80

TTT CTA CCC GAG GAT GCC ATT CTT AAA GGC GGT CAT TAT GAC AAC CAA 288
 Phe Leu Pro Glu Asp Ala Ile Leu Lys Gly Gly His Tyr Asp Asn Gln
 85 90 95

CTG CAA AAT GGC ATC AAG CGA GTA AAA GAG TTC CTT GAA TCA TCG CCG 336
 Leu Gln Asn Gly Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro
 100 105 110

AAT ACA CAA TGG GAA TTG CCG GCG TTC ATG GCA GTA ATG CAT TTC TCT 384
 Asn Thr Gln Trp Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser
 115 120 125

34

TTA ACC GCC GAT CGT ATC GAT GAT GAT ATT TTG AAA GTG ATT GTT GAT	432
Leu Thr Ala Asp Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp	
130 135 140	
TCR ATG AAT CAT CAT GGT GAT GCC CGT AGC AAG TTG CGT GAA GAA TTA	480
Ser Met Asn His His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu	
145 150 155 160	
GCT GAG CTT ACC GCC GAA TTA AAG ATT TAT TCA GTT ATT CAA GCC GAA	528
Ala Glu Leu Thr Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu	
165 170 175	
ATT AAT AAG CAT CTG TCT AGT AGT GGC ACC ATA AAT ATC CAT GAT AAA	576
Ile Asn Lys His Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys	
180 185 190	
TOC ATT AAT CTC ATG GAT AAA AAT TTA TAT GGT TAT ACA GAT GAA GAG	624
Ser Ile Asn Leu Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu	
195 200 205	
ATT TTT AAA GCC AGC GCA GAG TAC AAA ATT CTC GAG AAA ATG CCT CAA	672
Ile Phe Lys Ala Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln	
210 215 220	
ACC ACC ATT CAG GTG GAT GGG AGC GAG AAA AAA ATA GTC TCG ATA AAG	720
Thr Thr Ile Gln Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys	
225 230 235 240	

35

GAC TTT CTT GGA AGT GAG AAT AAA AGA ACC GGG GCG TTG GGT AAT CTG 768
 Asp Phe Leu Gly Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu

245

250

255

AAA AAC TCA TAC TCT TAT AAT AAA GAT AAT AAT GAA TTA TCT CAC TTT 816
 Lys Asn Ser Tyr Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe

260

265

270

GCC ACC ACC TGC TCG GAT AAG TCC AGG CCG CTC AAC GAC TTG GTT AGC 864
 Ala Thr Thr Cys Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser

275

280

285

CAA AAA ACA ACT CAG CTG TCT GAT ATT ACA TCA CGT TTT AAT TCA GCT 912
 Gln Lys Thr Thr Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala

290

295

300

ATT GAA GCA CTG AAC CGT TTC ATT CAG AAA TAT GAT TCA CTG ATG CAA 960
 Ile Glu Ala Leu Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln

305

310

315

320

CGT CTG CTA GAT GAC ACG TCT GGT AAA TGACACGAGG TAATTATGTA 1007
 Arg Leu Leu Asp Asp Thr Ser Gly Lys

325

AGTCGAC

1014

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 329 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Gly Ile Pro Gly Ile Arg Ala Tyr Glu Gln Asn Pro Gln His Phe Ile
1 5 10 15

Glu Asp Leu Glu Lys Val Arg Val Glu Gln Leu Thr Gly His Gly Ser
20 25 30

Ser Val Leu Glu Glu Leu Val Gln Leu Val Lys Asp Lys Asn Ile Asp
35 40 45

Ile Ser Ile Lys Tyr Asp Pro Arg Lys Asp Ser Glu Val Phe Ala Asn
50 55 60

Arg Val Ile Thr Asp Asp Ile Glu Leu Leu Lys Lys Ile Leu Ala Tyr
65 70 75 80

Phe Leu Pro Glu Asp Ala Ile Leu Lys Gly Gly His Tyr Asp Asn Gln
85 90 95

37

Leu Gln Asn Gly Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro
100 105 110

Asn Thr Gln Trp Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser
115 120 125

Leu Thr Ala Asp Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp
130 135 140

Ser Met Asn His His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu
145 150 155 160

Ala Glu Leu Thr Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu
165 170 175

Ile Asn Lys His Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys
180 185 190

Ser Ile Asn Leu Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu
195 200 205

Ile Phe Lys Ala Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln
210 215 220

Thr Thr Ile Gln Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys
225 230 235 240

Asp Phe Leu Gly Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu
245 250 255

38

Lys Asn Ser Tyr Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe
260 265 270

Ala Thr Thr Cys Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser
275 280 285

Gln Lys Thr Thr Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala
290 295 300

Ile Glu Ala Leu Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln
305 310 315 320

Arg Leu Leu Asp Asp Thr Ser Gly Lys
325

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GATCGAATTC ATTAGAGCCT ACGAACAA

28

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGATCGTCGA CTTACATAAT TACCTCGTGT CA

32

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

41

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7.

GATCCAGCTC GGCAGATTTA ACTGCAAGCA CC

31

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

CAGGTCGAGC TCGTCGACGG TTAGGCTCAA AGTAG

35

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

42

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(iii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

TGGTACGCTT ACTCTTGGCG GCTAT

25

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 541 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 2..454

(ix) FEATURE:

(A) NAME/KEY: misc_recomb

(B) LOCATION: 1..6

(ix) FEATURE:

(A) NAME/KEY: misc_recomb

(B) LOCATION: 536..541

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```
G AGC TCG GCA GAT TTA ACT GCA AGC ACC ACT GCA ACG GCA ACT GTT      46
Ser Ser Ala Asp Leu Thr Ala Ser Thr Thr Ala Thr Ala Thr Leu
1           5           10           15
```

```
GTT GAA CCA GCC CGC ATC ACT ATT ACA TAT AAG GAA GGC GGT CCA ATT      94
Val Glu Pro Ala Arg Ile Thr Ile Thr Tyr Lys Glu Gly Ala Pro Ile
20           25           30
```



44

ACA ATT ATG GAC AAT GGA AAC ATC GAT ACA GAA TTA CTT GTT GGT ACG 142
 Thr Ile Met Asp Asn Gly Asn Ile Asp Thr Glu Leu Leu Val Gly Thr

35

40

45

CTT ACT CTT GGC GGC TAT AAA ACA GGA ACC ACT AGC ACA TCT GTT AAC 190
 Leu Thr Leu Gly Gly Tyr Lys Thr Gly Thr Thr Ser Thr Ser Val Asn

50

55

60

TTT ACA GAT GCC GCG GGT GAT CCC ATG TAC TTA ACA TTT ACT TCT CAG 238
 Phe Thr Asp Ala Ala Gly Asp Pro Met Tyr Leu Thr Phe Thr Ser Gln

65

70

75

GAT GGA AAT AAC CAC CAA TTC ACT ACA AAA GTG ATT GGC AAG GAT TCT 286
 Asp Gly Asn Asn His Gln Phe Thr Thr Lys Val Ile Gly Lys Asp Ser

80

85

90

95

AGA GAT TTT GAT ATC TCT CCT AAG GTA AAC GGT GAG AAC CTT GTG GGG 334
 Arg Asp Phe Asp Ile Ser Pro Lys Val Asn Gly Glu Asn Leu Val Gly

100

105

110

GAT GAC GTC GTC TTG GCT ACG GGC AGC CAG GAT TTC TTT GTT CGC TCA 382
 Asp Asp Val Val Leu Ala Thr Gly Ser Gln Asp Phe Phe Val Arg Ser

115

120

125

ATT GGT TCC AAA GGC GGT AAA CTT GCA GCA GGT AAA TAC ACT GAT GCT 430
 Ile Gly Ser Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr Asp Ala

130

135

140

GTA ACC GTA ACC GTA TCT AAC CAA TAATCCATAT AGATAATAGA TAAAGGAGGG 484

Val Thr Val Thr Val Ser Asn Gln

145

150

CTATTATGCC CTCCTTTAAT APTTATGAAT TATCCTACTT TGAGCCTAAC CCTCGAC 541

46

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ser	Ser	Ala	Asp	Leu	Thr	Ala	Ser	Thr	Thr	Ala	Thr	Ala	Thr	Leu	Val
1				5				10						15	
Glu	Pro	Ala	Arg	Ile	Thr	Ile	Thr	Tyr	Lys	Glu	Gly	Ala	Pro	Ile	Thr
			20					25					30		
Ile	Met	Asp	Asn	Gly	Asn	Ile	Asp	Thr	Glu	Leu	Leu	Val	Gly	Thr	Leu
		35					40						45		
Thr	Leu	Gly	Gly	Tyr	Lys	Thr	Gly	Thr	Thr	Ser	Thr	Ser	Val	Asn	Phe
		50					55					60			
Thr	Asp	Ala	Ala	Gly	Asp	Pro	Met	Tyr	Leu	Thr	Phe	Thr	Ser	Gln	Asp
	65					70				75				80	
Gly	Asn	Asn	His	Gln	Phe	Thr	Thr	Lys	Val	Ile	Gly	Lys	Asp	Ser	Arg
			85						90					95	

47

Asp Phe Asp Ile Ser Pro Lys Val Asn Gly Glu Asn Leu Val Gly Asp
100 105 110

Asp Val Val Leu Ala Thr Gly Ser Gln Asp Phe Phe Val Arg Ser Ile
115 120 125

Gly Ser Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr Asp Ala Val
130 135 140

Thr Val Thr Val Ser Asn Gln
145 150

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Yersinia pestis

48

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TTGCTCTGGG AATCCGAACA TAAATCGGTT CAGTGGCC

38

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GGCCTATTCC TCGCTAGCAA TGTTTAACC

29

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 base pairs

49

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(x) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATCGTTAAAC ATTGCTAGCG AGGAATACGC C

31

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

50

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

GATAGATCTG TCGACTGAAC CTATTATATT GCTTCGCGC

39

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1462 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 8..1447

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GAGCTCG	GCA	GAT	TTA	ACT	GCA	AGC	ACC	ACT	GCA	ACG	GCA	ACT	CTT	GTT	45	
Ala Asp Leu Thr Ala Ser Thr Thr Ala Thr Ala Thr Leu Val																
1					5									10		
GAA	CCA	GCC	CGC	ATC	ACT	CTT	ACA	TAT	AAG	GAA	GGC	GCT	CCA	ATT	ACA	97
Glu Pro Ala Arg Ile Thr Leu Thr Tyr Lys Glu Gly Ala Pro Ile Thr																
15					20					25				30		
ATT	ATG	GAC	AAT	GGA	AAC	ATC	GAT	ACA	GAA	TTA	CTT	GTT	GGT	ACG	CTT	145
Ile Met Asp Asn Gly Asn Ile Asp Thr Glu Leu Leu Val Gly Thr Leu																
					35					40				45		
ACT	CTT	GGC	GGC	TAT	AAA	ACA	GGA	ACC	ACT	AGC	ACA	TCT	GTT	AAC	TTT	193
Thr Leu Gly Gly Tyr Lys Thr Gly Thr Thr Ser Thr Ser Val Asn Phe																
		50						55					60			
ACA	GAT	GCC	GCG	GGT	GAT	CCC	ATG	TAC	TTA	ACA	TTT	ACT	TCT	CAG	GAT	241
Thr Asp Ala Ala Gly Asp Pro Met Tyr Leu Thr Phe Thr Ser Gln Asp																
		65					70						75			
GGA	AAT	AAC	CAC	CAA	TTC	ACT	ACA	AAA	GTC	ATT	GGC	AAG	GAT	TCT	AGA	289
Gly Asn Asn His Gln Phe Thr Thr Lys Val Ile Gly Lys Asp Ser Arg																
	80						85						90			

52

GAT TTT GAT ATC TCT CCT AAG GTA AAC GGT GAG AAC CTT GTG GGG GAT 337
 Asp Phe Asp Ile Ser Pro Lys Val Asn Gly Glu Asn Leu Val Gly Asp
 95 100 105 110

GAC GTC GTC TTG GCT ACG GGC AGC CAG GAT TTC TTT GTT CGC TCA ATT 389
 Asp Val Val Leu Ala Thr Gly Ser Gln Asp Phe Phe Val Arg Ser Ile
 115 120 125

GGT TCC AAA GGC GGT AAA CTT GCA GCA GGT AAA TAC ACT GAT GCT GTA 433
 Gly Ser Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr Asp Ala Val
 130 135 140

ACC GTA ACC GTA TCT AAC CAA GGA TCC ATC GAA GGT CGT ATT AGA GCC 481
 Thr Val Thr Val Ser Asn Gln Gly Ser Ile Glu Gly Arg Ile Arg Ala
 145 150 155

TAC GAA CAA AAC CCA CAA CAT TTT ATT GAG GAT CTA GAA AAA GTT AGG 529
 Tyr Glu Gln Asn Pro Gln His Phe Ile Glu Asp Leu Glu Lys Val Arg
 160 165 170

GTG GAA CAA CTT ACT GGT CAT GGT TCT TCA GTT TTA GAA GAA TTG GTT 577
 Val Glu Gln Leu Thr Gly His Gly Ser Ser Val Leu Glu Glu Leu Val
 175 180 185 190

CAG TTA GTC AAA GAT AAA AAT ATA GAT ATT TCC ATT AAA TAT GAT CCC 625
 Gln Leu Val Lys Asp Lys Asn Ile Asp Ile Ser Ile Lys Tyr Asp Pro
 195 200 205

AGA AAA GAT TCG GAG GTT TTT GCC AAT AGA GTA ATT ACT GAT GAT ATC 673
 Arg Lys Asp Ser Glu Val Phe Ala Asn Arg Val Ile Thr Asp Asp Ile
 210 215 220

GAA TTG CTC AAG AAA ATC CTA GCT TAT TTT CTA CCC GAG GAT GCC ATT 721
 Glu Leu Leu Lys Lys Ile Leu Ala Tyr Phe Leu Pro Glu Asp Ala Ile
 225 230 235

CTT AAA GGC GGT CAT TAT GAC AAC CAA CTG CAA AAT GGC ATC AAG CGA 769
 Leu Lys Gly Gly His Tyr Asp Asn Gln Leu Gln Asn Gly Ile Lys Arg
 240 245 250

GTA AAA GAG TTC CTT GAA TCA TCG CCG AAT ACA CAA TGG GAA TTG CCG 817
 Val Lys Glu Phe Leu Glu Ser Ser Pro Asn Thr Gln Trp Glu Leu Arg
 255 260 265 270

GCG TTC ATG GCA GTA ATG CAT TTC TCT TTA ACC GCC GAT CGT ATC GAT 865
 Ala Phe Met Ala Val Met His Phe Ser Leu Thr Ala Asp Arg Ile Asp
 275 280 285

GAT GAT ATT TTG AAA GTG ATT GTT GAT TCA ATG AAT CAT CAT GGT GAT 913
 Asp Asp Ile Leu Lys Val Ile Val Asp Ser Met Asn His His Gly Asp
 290 295 300

GCC CGT AGC AAG TTG CGT GAA GAA TTA GCT GAG CTT ACC GCC GAA TTA 961
 Ala Arg Ser Lys Leu Arg Glu Glu Leu Ala Glu Leu Thr Ala Glu Leu
 305 310 315

AAG ATT TAT TCA GTT ATT CAA GCC GAA ATT AAT AAG CAT CTC TCT AGT 1009
 Lys Ile Tyr Ser Val Ile Gln Ala Glu Ile Asn Lys His Leu Ser Ser

320

325

330

AAT GGC ACC ATA AAT ATC CAT GAT AAA TCC ATT AAT CTC ATG GAT AAA 1087
 Ser Gly Thr Ile Asn Ile His Asp Lys Ser Ile Asn Leu Met Asp Lys

335

340

345

350

AAT TTA TAT GGT TAT ACA GAT GAA GAG ATT TTT AAA GCC AGC GCA GAG 1106
 Asn Leu Tyr Gly Tyr Thr Asp Glu Glu Ile Phe Lys Ala Ser Ala Glu

355

360

365

TAC AAA ATT CTC GAG AAA ATG CCT CAA ACC ACC ATT CAG GTG GAT GGG 1153
 Tyr Lys Ile Leu Glu Lys Met Pro Gln Thr Thr Ile Gln Val Asp Gly

370

375

380

AGC GAG AAA AAA ATA GTC TCG ATA AAG GAC TTT CTT GGA AGT GAG AAT 1201
 Ser Glu Lys Lys Ile Val Ser Ile Lys Asp Phe Leu Gly Ser Glu Asn

385

390

395

AAA AGA ACC GGG GCG TTG GGT AAT CTG AAA AAC TCA TAC TCT TAT AAT 1249
 Lys Arg Thr Gly Ala Leu Gly Asn Leu Lys Asn Ser Tyr Ser Tyr Asn

400

405

410

AAA GAT AAT AAT GAA TTA TCT CAC TTT GCC ACC ACC TGC TCG GAT AAG 1297
 Lys Asp Asn Asn Glu Leu Ser His Phe Ala Thr Thr Cys Ser Asp Lys

415

420

425

430

55

TCC AGG CCG CTC AAC GAC TTG GTT AGC CAA AAA ACA ACT CAG CTG TCT 1345
 Ser Arg Pro Leu Asn Asp Leu Val Ser Gln Lys Thr Thr Gln Leu Ser

435

440

445

GAT ATT ACA TCA CGT TTT AAT TCA GCT ATT CAA GCA CTG AAC CGT TTT 1390
 Asp Ile Thr Ser Arg Phe Asn Ser Ala Ile Glu Ala Leu Asn Arg Phe

450

455

460

ATT CAG AAA TAT GAT TCA GTG ATG CAA CGT CTG CTA GAT GAC ACG TCT 1441
 Ile Gln Lys Tyr Asp Ser Val Met Gln Arg Leu Leu Asp Asp Thr Ser

465

470

475

GGT AAA TGACACTAGA AGCTT

1482

Gly Lys

480

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 480 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ala Asp Leu Thr Ala Ser Thr Thr Ala Thr Ala Thr Leu Val Glu Pro

1

5

10

15

56

Ala Arg Ile Thr Leu Thr Tyr Lys Glu Gly Ala Pro Ile Thr Ile Met

20

25

30

Asp Asn Gly Asn Ile Asp Thr Glu Leu Leu Val Gly Thr Leu Thr Leu

35

40

45

Gly Gly Tyr Lys Thr Gly Thr Thr Ser Thr Ser Val Asn Phe Thr Asp

50

55

60

Ala Ala Gly Asp Pro Met Tyr Leu Thr Phe Thr Ser Gln Asp Gly Asn

65

70

75

80

Asn His Gln Phe Thr Thr Lys Val Ile Gly Lys Asp Ser Arg Asp Phe

85

90

95

Asp Ile Ser Pro Lys Val Asn Gly Glu Asn Leu Val Gly Asp Asp Val

100

105

110

Val Leu Ala Thr Gly Ser Gln Asp Phe Phe Val Arg Ser Ile Gly Ser

115

120

125

Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr Asp Ala Val Thr Val

130

135

140

Thr Val Ser Asn Gln Gly Ser Ile Glu Gly Arg Ile Arg Ala Tyr Glu

145

150

155

160

Gln Asn Pro Gln His Phe Ile Glu Asp Leu Glu Lys Val Arg Val Glu

165

170

175

57

Gln Leu Thr Gly His Gly Ser Ser Val Leu Glu Glu Leu Val Gln Leu
180 185 190

Val Lys Asp Lys Asn Ile Asp Ile Ser Ile Lys Tyr Asp Pro Arg Lys
195 200 205

Asp Ser Glu Val Phe Ala Asn Arg Val Ile Thr Asp Asp Ile Glu Leu
210 215 220

Leu Lys Lys Ile Leu Ala Tyr Phe Leu Pro Glu Asp Ala Ile Leu Lys
225 230 235 240

Gly Gly His Tyr Asp Asn Gln Leu Gln Asn Gly Ile Lys Arg Val Lys
245 250 255

Glu Phe Leu Glu Ser Ser Pro Asn Thr Gln Trp Glu Leu Arg Ala Phe
260 265 270

Met Ala Val Met His Phe Ser Leu Thr Ala Asp Arg Ile Asp Asp Asp
275 280 285

Ile Leu Lys Val Ile Val Asp Ser Met Asn His His Gly Asp Ala Arg
290 295 300

Ser Lys Leu Arg Glu Glu Leu Ala Glu Leu Thr Ala Glu Leu Lys Ile
305 310 315 320

Tyr Ser Val Ile Gln Ala Glu Ile Asn Lys His Leu Ser Ser Ser Gly
325 330 335

58

Thr Ile Asn Ile His Asp Lys Ser Ile Asn Leu Met Asp Lys Asn Leu
340 345 350

Tyr Gly Tyr Thr Asp Glu Glu Ile Phe Lys Ala Ser Ala Glu Tyr Lys
355 360 365

Ile Leu Glu Lys Met Pro Gln Thr Thr Ile Gln Val Asp Gly Ser Glu
370 375 380

Lys Lys Ile Val Ser Ile Lys Asp Phe Leu Gly Ser Glu Asn Lys Arg
385 390 395 400

Thr Gly Ala Leu Gly Asn Leu Lys Asn Ser Tyr Ser Tyr Asn Lys Asp
405 410 415

Asn Asn Glu Leu Ser His Phe Ala Thr Thr Cys Ser Asp Lys Ser Arg
420 425 430

Pro Leu Asn Asp Leu Val Ser Gln Lys Thr Thr Gln Leu Ser Asp Ile
435 440 445

Thr Ser Arg Phe Asn Ser Ala Ile Glu Ala Leu Asn Arg Phe Ile Gln
450 455 460

Lys Tyr Asp Ser Val Met Gln Arg Leu Leu Asp Asp Thr Ser Gly Lys
465 470 475 480

59

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ATAAGACTGT GCTAGCTAGA GGTAATATAT G

31

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

60

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GATGGATCCT TGGTTAGATA CGGTTACG

28

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 547 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 29..538

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATAAGACTGT GCTAGCTAGA GGTAAATAT ATG AAA AAA ATC AGT TCC GTT ATC	92
Met Lys Lys Ile Ser Ser Val Ile	
1 5	
GCC ATT GCA TTA TTT GCA ACT ATT GCA ACT GCT AAT GCG GCA GAT TTA	100
Ala Ile Ala Leu Phe Gly Thr Ile Ala Thr Ala Asn Ala Ala Asp Leu	
10 15 20	
ACT GCA AGC ACC ACT GCA ACG GCA ACT CTT GTT GAA CCA GCC CGC ATC	148
Thr Ala Ser Thr Thr Ala Thr Ala Thr Leu Val Glu Pro Ala Arg Ile	
25 30 35 40	
ACT CTT ACA TAT AAG GAA GCC GCT CCA ATT ACA ATT ATG GAC AAT GGA	196
Thr Leu Thr Tyr Lys Glu Gly Ala Pro Ile Thr Ile Met Asp Asn Gly	
45 50 55	
AAC ATC GAT ACA GAA TTA CTT GTT GGT ACG CTT ACT CTT GGC GGC TAT	244
Asn Ile Asp Thr Glu Leu Leu Val Gly Thr Leu Thr Leu Gly Gly Tyr	
60 65 70	
AAA ACA GGA ACC ACT AGC ACA TCT GTT AAC TTT ACA GAT GCC GCG GGT	292
Lys Thr Gly Thr Thr Ser Thr Ser Val Asn Phe Thr Asp Ala Ala Gly	
75 80 85	
GAT CCC ATG TAC TTA ACA TTT ACT TCT CAG GAT GGA AAT AAC CAC CAA	340
Asp Pro Met Tyr Leu Thr Phe Thr Ser Gln Asp Gly Asn Asn His Gln	
90 95 100	

62

TTC ACT ACA AAA GTG ATT GGC AAG GAT TCT AGA GAT TTT GAT ATC TCT 388

Phe Thr Thr Lys Val Ile Gly Lys Asp Ser Arg Asp Phe Asp Ile Ser

105

110

115

120

CCT AAG GTA AAC GGT GAG AAC CTT GTG GGG GAT GAC GTC GTC TTG GCT 436

Pro Lys Val Asn Gly Glu Asn Leu Val Gly Asp Asp Val Val Leu Ala

125

130

135

ACG GGC AGC CAG GAT TTC TTT GTT CGC TCA ATT GGT TCC AAA GGC GGT 484

Thr Gly Ser Gln Asp Phe Phe Val Arg Ser Ile Gly Ser Lys Gly Gly

140

145

150

AAA CTT GCA GCA GGT AAA TAC ACT GAT GCT GTA ACC GTA ACC GTA TCT 532

Lys Leu Ala Ala Gly Lys Tyr Thr Asp Ala Val Thr Val Thr Val Ser

155

160

165

AAC CAA GGATCCATC

547

Asn Gln

170

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 170 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

63

(xi) SEQUENCE DESCRIPTION: SEQ ID NO.21:

Met	Lys	Lys	Ile	Ser	Ser	Val	Ile	Ala	Ile	Ala	Leu	Phe	Gly	Thr	Ile
1				5					10					15	
Ala	Thr	Ala	Asn	Ala	Ala	Asp	Leu	Thr	Ala	Ser	Thr	Thr	Ala	Thr	Ala
			20					25					30		
Thr	Leu	Val	Glu	Pro	Ala	Arg	Ile	Thr	Leu	Thr	Tyr	Lys	Glu	Gly	Ala
		35					40					45			
Pro	Ile	Thr	Ile	Met	Asp	Asn	Gly	Asn	Ile	Asp	Thr	Glu	Leu	Leu	Val
		50				55				60					
Gly	Thr	Leu	Thr	Leu	Gly	Gly	Tyr	Lys	Thr	Gly	Thr	Thr	Ser	Thr	Ser
	65				70					75				80	
Val	Asn	Phe	Thr	Asp	Ala	Ala	Gly	Asp	Pro	Met	Tyr	Leu	Thr	Phe	Thr
			85						90					95	
Ser	Gln	Asp	Gly	Asn	Asn	His	Gln	Phe	Thr	Thr	Lys	Val	Ile	Gly	Lys
			100					105					110		
Asp	Ser	Arg	Asp	Phe	Asp	Ile	Ser	Pro	Lys	Val	Asn	Gly	Glu	Asn	Leu
		115					120					125			
Val	Gly	Asp	Asp	Val	Val	Leu	Ala	Thr	Gly	Ser	Gln	Asp	Phe	Phe	Val
	130					135						140			

64

Arg Ser Ile Gly Ser Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr
145 150 155 160

Asp Ala Val Thr Val Thr Val Ser Asn Gln
165 170

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1530 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(A) ORGANISM: *Yersinia pestis*

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 13..1515

65

(x) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TAGAGGTAAT AT ATG AAA AAA ATC AGT TCC GTT ATC GCC ATT GCA TTA	48
Met Lys Lys Ile Ser Ser Val Ile Ala Ile Ala Leu	
1 5 10	
TTT GGA ACT ATT GCA ACT GCT AAT GCG GCA GAT TTA ACT GCA AGC ACC	96
Phe Gly Thr Ile Ala Thr Ala Asn Ala Ala Asp Leu Thr Ala Ser Thr	
15 20 25	
ACT GCA ACG GCA ACT CTT GTT GAA CCA GCC CGC ATC ACT CTT ACA TAT	144
Thr Ala Thr Ala Thr Leu Val Glu Pro Ala Arg Ile Thr Leu Thr Tyr	
30 35 40	
AAG GAA GGC GCT CCA ATT ACA ATT ATG GAC AAT GGA AAC ATC GAT ACA	192
Lys Glu Gly Ala Pro Ile Thr Ile Met Asp Asn Gly Asn Ile Asp Thr	
45 50 55 60	
GAA TTA CTT GTT GGT ACG CTT ACT CTT GGC GGC TAT AAA ACA GGA ACC	240
Glu Leu Leu Val Gly Thr Leu Thr Leu Gly Gly Tyr Lys Thr Gly Thr	
65 70 75	
ACT AGC ACA TCT GTT AAC TTT ACA GAT GCC GCG GGT GAT CCC ATG TAC	288
Thr Ser Thr Ser Val Asn Phe Thr Asp Ala Ala Gly Asp Pro Met Tyr	
80 85 90	
TTA ACA TTT ACT TCT CAG GAT GGA AAT AAC CAC CAA TTC ACT ACA AAA	336
Leu Thr Phe Thr Ser Gln Asp Gly Asn Asn His Gln Phe Thr Thr Lys	
95 100 105	

66

GTG ATT GGC AAG GAT TCT AGA GAT TTT GAT ATC TCT CCT AAG GTA AAC 384
Val Ile Gly Lys Asp Ser Arg Asp Phe Asp Ile Ser Pro Lys Val Asn

110

115

120

GGT GAG AAC CTT GTG GGG GAT GAC GTC GTC TTG GCT ACG GGC ACC CAG 432
Gly Glu Asn Leu Val Gly Asp Asp Val Val Leu Ala Thr Gly Ser Gln

125

130

135

140

GAT TTC TTT GTT CGC TCA ATT GGT TCC AAA GGC GGT AAA CTT GCA GCA 480
Asp Phe Phe Val Arg Ser Ile Gly Ser Lys Gly Gly Lys Leu Ala Ala

145

150

155

GGT AAA TAC ACT GAT GCT GTA ACC GTA ACC GTA TCT AAC CAA GGA TCC 528
Gly Lys Tyr Thr Asp Ala Val Thr Val Thr Val Ser Asn Gln Gly Ser

160

165

170

ATC GAA GGT CGT ATT AGA GCC TAC GAA CAA AAC CCA CAA CAT TTT ATT 576
Ile Glu Gly Arg Ile Arg Ala Tyr Glu Gln Asn Pro Gln His Phe Ile

175

180

185

GAG GAT CTA GAA AAA GTT AGG GTG GAA CAA CTT ACT GGT CAT GGT TCT 624
Glu Asp Leu Glu Lys Val Arg Val Glu Gln Leu Thr Gly His Gly Ser

190

195

200

TCA GTT TTA GAA GAA TTG GTT CAG TTA GTC AAA GAT AAA AAT ATA GAT 672
Ser Val Leu Glu Glu Leu Val Gln Leu Val Lys Asp Lys Asn Ile Asp

205

210

215

220

ATT TCC ATT AAA TAT GAT CCC AGA AAA GAT TCG GAG GTT TTT GCC AAT 720
Ile Ser Ile Lys Tyr Asp Pro Arg Lys Asp Ser Glu Val Phe Ala Asn

225

230

235

AGA GTA ATT ACT GAT GAT ATC GAA TTG CTC AAG AAA ATC CTA GCT TAT 768

Arg Val Ile Thr Asp Asp Ile Glu Leu Leu Lys Lys Ile Leu Ala Tyr

240

245

250

TTT CTA CCC GAG GAT GCC ATT CTT AAA GGC GGT CAT TAT GAC AAC CAA 816

Phe Leu Pro Glu Asp Ala Ile Leu Lys Gly Gly His Tyr Asp Asn Gln

255

260

265

CTG CAA AAT GGC ATC AAG CGA GTA AAA GAG TTC CTT GAA TCA TCG CCG 864

Leu Gln Asn Gly Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro

270

275

280

AAT ACA CAA TGG GAA TTG CGG GCG TTC ATG GCA GTA ATG CAT TTC TCT 912

Asn Thr Gln Trp Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser

285

290

295

300

TTA ACC GCC GAT CGT ATC GAT GAT GAT ATT TTG AAA GTG ATT GTT GAT 960

Leu Thr Ala Asp Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp

305

310

315

TCA ATG AAT CAT CAT GGT GAT GCC CGT AGC AAG TTG CGT GAA GAA TTA 1008

Ser Met Asn His His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu

320

325

330

GCT GAG CTT ACC GCC GAA TTA AAG ATT TAT TCA GTT ATT CAA GCC GAA 1056

Ala Glu Leu Thr Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu

335

340

345

ATT AAT AAG CAT CTG TCT AGT AGT GGC ACC ATA AAT ATC CAT GAT AAA 1184
 Ile Asn Lys His Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys
 350 355 360

TCC ATT AAT CTC ATG GAT AAA AAT TTA TAT GGT TAT ACA GAT GAA GAG 1182
 Ser Ile Asn Leu Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu
 365 370 375 380

ATT TTT AAA GCC AGC CCA GAG TAC AAA ATT CTC GAG AAA ATG CCT CAA 1200
 Ile Phe Lys Ala Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln
 385 390 395

ACC ACC ATT CAG GTG GAT GGG AGC GAG AAA AAA ATA GTC TCG ATA AAG 1248
 Thr Thr Ile Gln Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys
 400 405 410

GAC TTT CTT GGA AGT GAG AAT AAA AEA ACC GGG GCG TTG GGT AAT CTG 1296
 Asp Phe Leu Gly Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu
 415 420 425

AAA AAC TCA TAC TCT TAT AAT AAA GAT AAT AAT GAA TTA TCT CAC TTT 1344
 Lys Asn Ser Tyr Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe
 430 435 440

GCC ACC ACC TGC TCG GAT AAG TCC AGG CCG CTC AAC GAC TTG GTT AGC 1392
 Ala Thr Thr Cys Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser
 445 450 455 460

69

CAA AAA ACA ACT CAG CTG TCT GAT ATT ACA TCA CGT TTT AAT TCA GCT 1440
 Gln Lys Thr Thr Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala
 465 470 475

ATT GAA GCA CTG AAC CTT TTC ATT CAG AAA TAT GAT TCA GTG ATG CAA 1488
 Ile Glu Ala Leu Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln
 480 485 490

CGT CTG CTA GAT GAC ACG TCT GGT AAA TGACACTAGA AGCTT 1530
 Arg Leu Leu Asp Asp Thr Ser Gly Lys
 495 500

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 501 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

Met Lys Lys Ile Ser Ser Val Ile Ala Ile Ala Leu Phe Gly Thr Ile
 1 5 10 15
 Ala Thr Ala Asn Ala Ala Asp Leu Thr Ala Ser Thr Thr Ala Thr Ala
 20 25 30

70

Thr Leu Val Glu Pro Ala Arg Ile Thr Leu Thr Tyr Lys Glu Gly Ala
35 40 45

Pro Ile Thr Ile Met Asp Asn Gly Asn Ile Asp Thr Glu Leu Leu Val
50 55 60

Gly Thr Leu Thr Leu Gly Gly Tyr Lys Thr Gly Thr Thr Ser Thr Ser
65 70 75 80

Val Asn Phe Thr Asp Ala Ala Gly Asp Pro Met Tyr Leu Thr Phe Thr
85 90 95

Ser Gln Asp Gly Asn Asn His Gln Phe Thr Thr Lys Val Ile Gly Lys
100 105 110

Asp Ser Arg Asp Phe Asp Ile Ser Pro Lys Val Asn Gly Glu Asn Leu
115 120 125

Val Gly Asp Asp Val Val Leu Ala Thr Gly Ser Gln Asp Phe Phe Val
130 135 140

Arg Ser Ile Gly Ser Lys Gly Gly Lys Leu Ala Ala Gly Lys Tyr Thr
145 150 155 160

Asp Ala Val Thr Val Thr Val Ser Asn Gln Gly Ser Ile Glu Gly Arg
165 170 175

Ile Arg Ala Tyr Glu Gln Asn Pro Gln His Phe Ile Glu Asp Leu Glu
180 185 190

Lys Val Arg Val Glu Gln Leu Thr Gly His Gly Ser Ser Val Leu Glu
195 200 205

Glu Leu Val Gln Leu Val Lys Asp Lys Asn Ile Asp Ile Ser Ile Lys
210 215 220

Tyr Asp Pro Arg Lys Asp Ser Glu Val Phe Ala Asn Arg Val Ile Thr
225 230 235 240

Asp Asp Ile Glu Leu Leu Lys Lys Ile Leu Ala Tyr Phe Leu Pro Glu
245 250 255

Asp Ala Ile Leu Lys Gly Gly His Tyr Asp Asn Gln Leu Gln Asn Gly
260 265 270

Ile Lys Arg Val Lys Glu Phe Leu Glu Ser Ser Pro Asn Thr Gln Trp
275 280 285

Glu Leu Arg Ala Phe Met Ala Val Met His Phe Ser Leu Thr Ala Asp
290 295 300

Arg Ile Asp Asp Asp Ile Leu Lys Val Ile Val Asp Ser Met Asn His
305 310 315 320

His Gly Asp Ala Arg Ser Lys Leu Arg Glu Glu Leu Ala Glu Leu Thr
325 330 335

Ala Glu Leu Lys Ile Tyr Ser Val Ile Gln Ala Glu Ile Asn Lys His
340 345 350

72

Leu Ser Ser Ser Gly Thr Ile Asn Ile His Asp Lys Ser Ile Asn Leu
 355 360 365

Met Asp Lys Asn Leu Tyr Gly Tyr Thr Asp Glu Glu Ile Phe Lys Ala
 370 375 380

Ser Ala Glu Tyr Lys Ile Leu Glu Lys Met Pro Gln Thr Thr Ile Gln
 385 390 395 400

Val Asp Gly Ser Glu Lys Lys Ile Val Ser Ile Lys Asp Phe Leu Gly
 405 410 415

Ser Glu Asn Lys Arg Thr Gly Ala Leu Gly Asn Leu Lys Asn Ser Tyr
 420 425 430

Ser Tyr Asn Lys Asp Asn Asn Glu Leu Ser His Phe Ala Thr Thr Cys
 435 440 445

Ser Asp Lys Ser Arg Pro Leu Asn Asp Leu Val Ser Gln Lys Thr Thr
 450 455 460

Gln Leu Ser Asp Ile Thr Ser Arg Phe Asn Ser Ala Ile Glu Ala Leu
 465 470 475 480

Asn Arg Phe Ile Gln Lys Tyr Asp Ser Val Met Gln Arg Leu Leu Asp
 485 490 495

Asp Thr Ser Gly Lys
 500

73

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 24 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(v) ORIGINAL SOURCE:

- (A) ORGANISM: Yersinia pestis

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GATCGAATTC GAGCTTACCA ACAA

24

As discussed previously in this application, the V and F1 antigen may be microencapsulated. The V and F1 antigens may be separately microencapsulated. The combined microencapsulated sub-units may be used for immunisation. The present inventors believe that the protection afforded by the combined microencapsulated sub-units is superior to that provided by existing plague vaccines and that there is an additive effect in combining the sub-units. The protective efficacy of the combined microencapsulated sub-units may further be enhanced by co-administering an adjuvant for example cholera toxin B sub-unit (CTB). Microencapsulation of the sub-unit vaccine prolongs release of the vaccine *in vivo*, permits oral, intra-nasal or inhalational delivery and gives scope for targetting.

The microencapsulation of the combined V and F1 sub-unit vaccine is described below. Also demonstrated is that this formulation is able to induce both mucosal and systemic immunity against plague.

The microencapsulation of sub-units was effected in PLA 2000 using a solvent evaporation technique.

Immunisation with the microencapsulated sub-units was carried out as follows.

Groups of 21 mice received a primary immunising dose of 25 μ g of either V antigen or F1 antigen, presented in microspheres resuspended in PBS for intra-peritoneal (i.p.) injection. Further groups of 21 mice received a combination of 25 μ g of each of the F1 and V antigens, presented in microspheres. A dose of 25 μ g of F1 was delivered in a total mass of 5.42mg of spheres, whilst 25 μ g of V was contained in 2.08mg of spheres. The required mass of microspheres was re-suspended in 100 μ l PBS per animal for injection. Animals were boosted with the respective antigen(s), as appropriate, on days 14 and 28.

Two further groups of 21 mice were primed and then boosted i.p. on days 14 and 28 with a combination of 10 μ g F1 and 10 μ g V, jointly incorporated in the aqueous phase of 25% (v/v) suspension of alhydrogel (Alhydrogel 1.3%, Superfos, Denmark) in PBS.

Selected groups of animals received in addition a dose of 10µg CTB (Sigma, Poole) incorporated into the delivery vehicle at each dosing point.

Control groups, each of 21 mice, received either alhydrogel only (100µl of 25% solution) or CTB only (10µg in 100µl PBS) or remained untreated.

In order to compare the protective efficacy of immunisation with combined sub-units, free or microencapsulated, against that provided by the Greer vaccine, (purchased from Greer laboratories) animals were challenged s.c. with virulent *Y.pestis*.

There was a 60% survival rate in Greer vaccinees against a challenge of 2×10^5 cfu *Y.pestis* (Fig. 1). By comparison, 80 % of the combined microencapsulated F1 + V (group 1) survived this challenge and there was 100% survival in group 2 (µV + µF1 + 10µg CTB). Thus the combined microencapsulated formulation was protective against virulent *Y. pestis* with no evidence of side effects.

In summary treatment groups were:

Group	Treatment
1.	25µg microencapsulated F1 (µ F1) + 25µg microencapsulated V (µV) i.p.
2.	25µg µ F1 + 25µg µV + 10µg CTB i.p.
3.	25µg µ F1 i.p.
4.	25µV i.p.
5.	25µg F1 + 25µg V in alhydrogel i.p.
6.	Greer vaccine 0.1ml i.m.

Micro-encapsulation may also be carried out with block co-polymers. In the following experiments, model protein antigen BSA was used. The preparation and characterisation of microspheres is as follows. Protein-loaded microspheres were prepared by an oil/water solvent evaporation method procedure previously described see R.L. Hunter and B. Bennet, *The J. Immunol.*, 133(6), 3167-3175 (1984), with some modifications. Polymer (poly-D-lactic acid): Resomer 206, Boehringer Ingelheim, Germany; 125 or 250mg) solution in acetone (22.5ml), containing model protein (antigen) BSA (at 15-25% theoretical loading level) and 0.11% w/v Pluronic L101 (or 0.09% w/v L121) available from Zeneca, probe sonicated for 10 seconds and then added to the aqueous phase (22.5ml), mixing at 100 rpm for 5 minutes and rotary evaporated until the organic solvent had been removed. The resulting colloid was washed and freeze-dried. Microspheres with an average diameter of $\sim 1\mu\text{m}$ (as determined by Malvern Mastersizer) and protein loadings ~ 0.5 -1.0% produced in this fabrication condition. External morphology of the resulting microspheres were analysed by scanning electron microscopy (SEM). Surface characteristics were defined in terms of zeta potential and hydrophobicity.

Hydrophobicity measurements: hydrophobicity of microspheres was quantified using hydrophobic interaction chromatography (HIC) as previously reported see H.O. Alpar and A.J. Almeida, *Eur. J. Pharm. Biopharm.*, 40, (4), 198-202 (1994). Microspheres were eluted from a series of agaroses which were modified with hydrophobic residues. The retention of microspheres in octyl agarose was used as an index of hydrophobicity.

Immunisation: A study was designed to establish the effects of differences in the type of microsphere and surface properties of the immune response. Female Balb/c mice (five per group) were injected i.m. with a single dose of BSA either encapsulated in Pluronic formulated microspheres or free in 100 μl alone, or suspended in the presence of surfactant. The control group of mice received the same amount of antigen encapsulated into microspheres containing PVA as emulsion stabiliser. Tail tip blood samples were removed periodically for 2 months. The serum from each sample was analysed for anti-BSA antibody using an enzyme-linked immunosorbent assay (ELISA). Results are represented graphically in Figure 2.

The presence of Pluronic L101 and L121 endows the surface with a much higher degree of hydrophobicity compared to PVA formulations (70% retained on octyl agarose column as opposed to 30% latex control 95% retained). A hydrophobic surface would facilitate macrophage interactions and subsequent uptake and would therefore be much more likely to mediate increased immune response. Figure 2 shows the effect of different BSA formulations in eliciting an immunoresponse. Batches formed by Pluronic L101 had a more enhanced effect on the plasma anti-BSA antibody titre than those formed by PVA. Batches formed by Pluronic L121 were slightly inferior to those of PVA microspheres in inducing good primary antibody response after delivery of only a small dose of antigen (1 μ g). The higher serum IgG level obtained with Pluronic L101 preparations as compared to other preparations is noted and may partly be due to the higher surface hydrophobicity.

it was also mentioned earlier in the present application that DNA encoding the whole or part of the F1 antigen and DNA encoding the whole or part of the V antigen could be used directly as a genetic vaccine. By way of example this may be carried out as follows.

1/ DNA encoding *Y.pestis* F1 and V is obtained by Polymerase Chain Reaction (PCR) amplification of specific regions of the *Y.pestis* genome, or by isolation of these genes from previously constructed plasmid clones e.g. for V exemplified sequence I.D. no 3.

2/ The F1 and V genes are cloned into mammalian expression vector plasmids such that the genes are situated downstream of a eukaryotic promoter. Suitable plasmids include pCMV3 (purchased from Clontech), in which the cytomegalovirus Immediate Early promoter is used. F1 and V may be cloned individually or in combination, and may be cloned as fusions with such genes as glutathione S-transferase, or eukaryotic signal sequences, which may stabilise the expressed protein and may facilitate export from mammalian cells.

3/ The recombinant plasmids are propagated in *Escherichia coli* and stocks are purified for transfection into a mammalian animal model and for immunisation of experimental animals by the intra-muscular or intra-dermal or inhalational routes.

Example 1: To construct a DNA vaccine expressing V antigen, the plasmid vector pCMV was digested with restriction enzyme *Not* 1 to remove the *lac* Z gene coding sequence. The digested plasmid was treated with Klenow enzyme to create blunt-ended vector DNA. An *ssp* 1 restriction fragment containing the coding sequence for a fusion protein of V antigen and glutathione-S transferase was isolated from recombinant plasmid pVG100 and ligated to the vector DNA. The V sequence used in this case is that described in exemplified Seq. ID No. 3. The recombinant plasmid was transformed into *E. coli* strain Nova Blue. Purified plasmid was inoculated into Balb/c mice by intra-muscular injection. Immunoglobulin responses to V antigen were detected in the serum of inoculated animals.

Example 2: To construct a DNA vaccine expressing F1 antigen, PCR primers were designed to amplify the complete *caf1* open reading frame. This encodes F1 and its signal peptide which directs export of the protein from the bacterial cell. The PCR primers had "tails" at their 5' ends which contained restriction enzyme recognition sites to allow directional insertion into a plasmid vector. The sequences of the PCR primers, 5'FAB2 and 3'FBAM, are given in exemplified Seq.ID. No. 18 and Seq.ID No. 19, respectively.

5'FAB2 and 3'FBAM were used to amplify a PCR fragment, the sequence of which is given in exemplified Seq.ID.no. 20. The PCR fragment was digested with restriction enzymes *Nhe* 1 and *Bam* *H*I and cloned into the plasmid pBKCMV which had been digested with the same enzymes. The resulting plasmid, pF1AB was transformed into *E. coli* Nova Blue and purified plasmid was used to inoculate Balb/c mice by intra-muscular injection. Immunoglobulin responses to F1 were found in inoculated animals.

Example 3: To construct a DNA vaccine expressing both F1 and V, the coding sequence for V was inserted into the DNA vaccine expressing F1, detailed in example 2. A linker region coding for 6 amino acids was positioned between the F1 and V coding sequences to allow each of the proteins to attain their conformational shape independently. The linker-V coding sequence was obtained by digesting the recombinant plasmid *placFV6* with *Bam HI* and *Hind III*. The linker-V DNA was ligated with the plasmid *pFIAB* which had also been digested with *Bam HI* and *Hind III*. The resulting plasmid, *pFVAB*, was transformed into cells of *E. coli* Nova Blue and stocks of plasmid were purified for further use. The nucleotide and derived amino acid sequence of the F1/V fusion are given in exemplified Seq.ID No. 22.

An example of a fusion protein comprising both F1 and V antigens is described below.

Enzymes and reagents.

Materials for the preparation of growth media were obtained from Oxoid Ltd or Difco Laboratories. All enzymes used in the manipulation of DNA were obtained from Boehringer Mannheim UK Ltd and used according to the manufacturer's instructions. All other chemicals and biochemicals were obtained from Sigma Chemical Co unless otherwise indicated. Monospecific rabbit polyclonal anti-V serum was supplied by Dr R Brubaker (Department of Microbiology, Michigan State University) and mouse anti-F1 IgA monoclonal antibody (Mab) F13G8-1 was obtained from the American Type Culture Collection.

Bacterial strains and cultivation.

Yersinia pestis GB was cultured aerobically at 28°C in a liquid medium (pH 6.8) containing 15 g proteose peptone, 2.5 g liver digest, 5 g yeast extract, 5 g NaCl per litre, supplemented with 80 ml of 0.25% haemin dissolved in 0.1M NaOH (Blood Base broth). *Escherichia coli* JM109 was cultured and stored as described by Sambrook et al (Sambrook J et al. 1989. Molecular cloning, a laboratory manual. Cold Spring Harbor Laboratory Press, New York).

Manipulation of DNA.

Chromosomal DNA was isolated from *Y. pestis* by the method of Marmur (Marmur, J. 1961. J Mol Biol 3: 206-218). The genes encoding F1 antigen (*cafI*) and V antigen (*icrV*) were amplified from *Y. pestis* DNA using PCR with 125 pmol of primers homologous to sequences from the 5' and 3' ends of the gene (Galyov, E E et al. 1990. FEBS Lett

277:230-232; Price S B et al. 1989. J Bacteriol 171: 5646-5653), although for *cafI* only the region encoding the mature F1 antigen was amplified. The sequences of the F1 5' primer (F/5'B: GATCGAGCTCGGCAGATTTAAGTCAAG CACC), the F1 3' primer (Flink/3'A: GCATGGATCCTTGGTTAGATACGGTTACGGT), the V 5' primer (Vlink/5'A: ATGGATCCATCGAAGGTCGTATTAGAGCCTACGAACAA), and the V 3' primer (VG/3'A: GCATAAGCTTCTA*GTGTCATTTACCAGACGT) also included 5' tails encoding the restriction sites *SacI*, *BamHI*, *BamHI* and *HindIII*, respectively. In addition, the nucleotide A* was altered from the published sequence of *lcrV* (Price S B et al. 1989. J Bacteriol 171: 5646-5653) to include an extra termination codon (TAA) in the amplified DNA. The tail of primer Vlink/5'A also included nucleotides encoding the factor Xa cleavage sequence Ile-Glu-Gly-Arg. The PCR primers were prepared with a DNA synthesiser (model 392; Applied Biosystems). DNA fragments were obtained after 30 cycles of amplification (95°C, 20 s; 45°C, 20 s; 72°C, 30 s; model 9600 GeneAmp PCR System; Perkin Elmer) and the fragments were purified. The *cafI* PCR product was digested with *SacI* and *BamHI*, ligated with suitably digested plasmid pUC18 and transformed into *E. coli* JM109 by electroporation. Subsequently, the *lcrV*-linker PCR product was digested with *BamHI* and *HindIII*, and ligated into the intermediate plasmid to form the recombinant plasmid *placFV6*. A colony containing *placFV6* was identified by PCR using 30-mer primers (5' nucleotides located at positions 54 and 794 (Price S B et al. 1989. J Bacteriol 171: 5646-5653) which amplified an internal segment of the *lcrV* gene. To confirm the nucleotide sequence of the cloned insert, sequencing reactions containing *placFV6* and primers designed from the *cafI* and *lcrV* genes were performed using an automated Taq polymerase cycle sequencing protocol with fluorescently labelled dideoxy nucleotides (CATALYST Molecular Biology Labstation; Applied Biosystems). The reaction products were analysed using an automated DNA sequencer (model 373A; Applied Biosystems).

The DNA sequence and derived amino acid sequence of the cloned fusion protein is shown in Example 1. The fusion protein consists of F1 and V antigens separated by a six-amino acid linker Gly-Ser-Ile-Glu-Gly-Arg. It is cloned downstream of the *lac* promoter and

in-frame with the vector-encoded LacZ' fragment. Thus, the complete fusion protein encodes nine additional amino acids at the N-terminus (Met-Thr-Met-Ile-Thr-Asn-Ser-Ser-Ser), and it accumulates in the cytoplasm.

Expression of the F1/V fusion protein in *E. coli*.

Cultures of *E. coli* JM109/placFV6 were grown in LB containing $100 \mu\text{gml}^{-1}$ ampicillin at 37°C until the absorbance (600nm) was 0.3. Isopropyl- β -D-thiogalactopyranoside (IPTG) was then added to the culture to a final concentration of 1 mM and growth was continued for a further 5 h. Whole cell lysates of the bacteria were prepared as described by Sambrook et al (Sambrook J et al. 1989, Molecular cloning, a laboratory manual, Cold Spring Harbor Laboratory Press, New York) and expression of the F1/V fusion protein was examined by SDS-PAGE on 10-15% gradient gels (Phastsystem, Pharmacia Biotech) and Western blotting (Sambrook J et al. 1989, Molecular cloning, a laboratory manual, Cold Spring Harbor Laboratory Press, New York). Western blots were probed with rabbit anti-V serum at a dilution of 1/4000 or Mab F13G8-1 at a dilution of 1/250, and protein bands were visualised with a colloidal gold labelled secondary antibody (Auroprobe BLplus, Cambio) or an anti-mouse IgA secondary antibody conjugated to horse radish peroxidase (Sigma).

A fusion protein with an approximate molecular weight of 54.2 kDa was detected in lysates of JM109/placFV6 by Western blotting with anti-V and anti-F1 sera. This product was not detected in control lysates of JM109/pUC18.

Electroporation into *Salmonella typhimurium* SL3261.

Plasmid DNA was extracted and purified from JM109/placFV6 using a Qiaprep kit (Qiagen) and electroporated into *S. typhimurium* strain LB5010 ($r^- m^+$). Subsequently, modified placFV6 was isolated and electroporated in *S. typhimurium* strain SL3261 (*aroA his*). For inoculation into mice, bacteria were grown in LB containing $100 \mu\text{gml}^{-1}$ ampicillin for 18 h

without shaking. After washing, the cells were resuspended in 10% glycerol in phosphate buffered saline (PBS) and stored at -70°C . The cell suspensions were defrosted and diluted in PBS as required prior to injection.

Immunisation with SL3261/placFV6.

Six week old female Balb/c mice, raised under specific pathogen-free conditions (Charles-River Laboratories, Margate, Kent, UK), were used in this study. A group of 19 mice received 0.1 ml immunising doses of approximately 5×10^6 cfu of SL3261/pFV6 on days 0 and 14 by the intravenous (iv) route. To retain placFV6 *in vivo*, mice were also injected subcutaneously (sc) with 50 μl ampicillin trihydrate suspension (150 mg ml^{-1} ; Penbritin injectable suspension POM; SmithKline Beecham Animal Health) for 5 days after each immunisation. In addition, groups of 15 mice were immunised iv on day 0 with a single 0.1 ml dose of approximately 5×10^6 cfu of SL3261 or intraperitoneally (ip) on days 0 and 14 with 0.1 ml of a mixture of 10 μg V and 10 μg F1 adsorbed to Alhydrogel. An untreated group of 10 age-matched mice were used as controls.

On day 7, five mice from the groups receiving SL3261/placFV6 or SL3261 were sacrificed and their spleens were removed. The organs were homogenised in 5 ml of PBS with a stomacher (Seward Medical Ltd) for 30 sec. The homogenates were serially diluted in PBS and inoculated on to L-agar or L-agar containing $100 \text{ } \mu\text{g ml}^{-1}$ ampicillin to determine the number of bacteria per spleen.

<i>S. typhimurium</i>	Actual dose	Average cfu per spleen \pm sem ^a		% recombinant
		L-agar	L-amp	
SL3261/placFV6	3.3×10^6 cfu	1030 ± 294	360 ± 135	37%
SL3261	1.6×10^7 cfu	$1.85 \times 10^7 \pm$ 3.57×10^6		

^a standard error of the mean

Measurement of serum antibody titre.

On day 42, blood was sampled from the tail vein of mice immunised with SL3261/placFV6 and pooled. The serum anti-V and anti-F1 titres were measured by a modified ELISA (Williamson, E D and R W Tilball, 1993, Vaccine 11:1253-1258). Briefly, V ($5 \mu\text{gml}^{-1}$ in PBS) or F1 ($2 \mu\text{gml}^{-1}$) were coated on to a microtitre plate and the test sera were serially diluted in duplicate on the plate. Bound antibody was detected using peroxidase labelled conjugates of anti-mouse polyvalent Ig. The titre of specific antibody was estimated as the maximum dilution of serum giving an absorbance reading greater than 0.1 units, after subtraction of the absorbance due to non-specific binding detected in the control sera. The serum antibody titre was also determined for all groups of mice prior to challenge with *Y. pestis*.

On day 42, the anti-V and anti-F1 titres of mice receiving SL3261/placFV6 were 1:5120 and 1:2560, respectively.

Challenge with *Y. pestis*.

On day 57, groups of 5 or 7 mice from the immunised and control groups were challenged subcutaneously with 0.1 ml aliquots of *Y. pestis* strain GB containing 7.36×10^2 or 7.36×10^4 cfu. Strain GB was isolated from a fatal human case of plague and has a median lethal dose (MLD) of < 1 cfu in Balb/c mice by the s.c. route (Russell, P et al. 1995. Vaccine 13: 1551-1556). The mice were observed for 14 days and, where appropriate, the time to death was recorded. A post-mortem was carried out on all animals where possible. To test for the presence of *Y. pestis*, samples of blood, liver and spleen were smeared on to Congo Red agar and incubated at 28°C for 48 h.

Group		Actual challenge dose - cfu	Survivors	MTD \pm sem* - hours
IV	SL3261/ placFV6	7.36×10^2	6/7	320
		7.36×10^4	6/7	124
	SL3261	7.36×10^2	0/5	97.4 ± 17.8
		7.36×10^4	0/5	97.6 ± 18.9
IP	F1 - V	7.36×10^2	7/7	-
		7.36×10^4	6/7	184
	Controls	7.36×10^2	0/5	116.8 ± 4.8
		7.36×10^4	0/5	63.6 ± 6.9

* standard error of the mean

CLAIMS

1. A method of protecting a human or animal body from the effects of infection with Y. pestis comprising administering to the body a vaccine including Yersinia pestis V antigen and Yersinia pestis F1 antigen or a protective epitopic part of each of these in a form other than whole Y. Pestis organisms.
2. A method as claimed in claim 1 wherein the antigens are administered in the form of a live vaccine.
3. A method as claimed in claim 2 wherein the live vaccine comprises human or animal gut colonising organisms that have been transformed using recombinant DNA to enable each organism to express one or both of V antigen and F1 antigen.
4. A method as claimed in claim 3 wherein the gut colonising organisms have been transformed with recombinant DNA such that they are enabled to express a fusion protein comprising both V and F1 antigen amino acid sequences or a protective epitopic part of each.
5. A method as claimed in claim 3 or 4 wherein the DNA comprises DNA of SEQ ID No 1 or SEQ ID No 3.
6. A method as claimed in claim 5 wherein the DNA is positioned in frame with a lacZ or nir β promoter.
7. A method as claimed in claim 3 or 4 wherein the DNA comprises DNA of SEQ ID No 10.
8. A method as claimed in any one of the preceding claims wherein the vaccine comprises isolated and/or purified recombinant V and F1 antigens.

9. A method as claimed in claim 8 wherein the antigens are provided in a pharmaceutically acceptable carrier.

10. A method as claimed in claim 9 wherein the carrier such as to produce an oil-in-water emulsion.

11. A method as claimed in any one of the preceding claims wherein the vaccine includes an adjuvant.

12. A method as claimed in any one of the preceding claims wherein the vaccine is administered such that it is enabled to induce local stimulation of the gut-associated lymphoid tissue (GALT) and, by trafficking of lymphocytes through the common mucosal immune system provide a secondary stimulation of the bronchial associated lymphoid tissue (BALT) such that a secretory IgA response is achieved at the respiratory mucosal surface.

13. A method as claimed in any one of the preceding claims wherein the vaccine is in the form of droplets or capsules.

14. A method as claimed in claim 13 wherein the capsules are liposomes or microcapsules effective in delivering the composition to the airways of an individual for the purposes of evoking mucosal immune response.

15. A vaccine composition comprising Yersinia pestis V antigen and Yersinia pestis F1 antigen or a protective epitopic part of each of these in a form other than whole Y. Pestis organisms.

16. A vaccine as claimed in claim 15 characterised in that it is a live vaccine.

17. A vaccine as claimed in claim 16 wherein the live vaccine comprises human or animal gut colonising organisms that have been transformed using recombinant DNA to enable them to express one or both of V antigen and F1 antigen.

18. A vaccine as claimed in claim 17 wherein the gut colonising organisms have been transformed with recombinant DNA such that they are enabled to express a fusion protein comprising both V and F1 antigen amino acid sequences or a protective epitopic part of each.
19. A vaccine as claimed in claim 17 or 18 wherein the DNA comprises DNA of SEQ ID No 1 or SEQ ID No 3.
20. A vaccine as claimed in claim 19 wherein the DNA is positioned in frame with a lacZ or nir β promoter.
21. A vaccine as claimed in claim 17 or 18 wherein the DNA comprises DNA of SEQ ID No 10.
22. A vaccine as claimed in any one of the preceding claims wherein the vaccine comprises isolated and/or purified recombinant V and F1 antigens.
23. A vaccine as claimed in claim 22 wherein the antigens are provided in a pharmaceutically acceptable carrier.
24. A vaccine as claimed in claim 23 wherein the carrier is such as to produce an oil-in-water emulsion.
25. A vaccine as claimed in any one of the preceding claims characterised in that it includes an adjuvant.
26. A vaccine as claimed in any one of preceding claims characterised in that it is in the form of droplets or capsules.

27. A vaccine as claimed in claim 26 wherein the capsules are liposomes or microcapsules effective in delivering the composition to the airways of an individual for the purposes of evoking mucosal immune response.
28. A vaccine as claimed in claim 26 wherein the capsules are block co-polymers.
29. A vaccine as claimed in claim 26 wherein the capsules comprise biodegradable polymers.
30. A vaccine as claimed in claim 29 wherein the biodegradable polymer is poly-lactic acid.
31. A vaccine as claimed in claim 30 further comprising glycollic acid.
32. A vaccine as claimed in claim 30 further comprising block co-polymer.
33. A vaccine according to either of claims 28 or 32 in which the block co-polymer contains the repeat unit $(POP-POE)_n$

34. A method as claimed in claim 5 wherein the DNA is positioned in frame with an in-vivo inducible promoter.
35. A method according to claim 34 wherein the in-vivo inducible promoter is selected from HtrA, nir β , OmpR, OmpC, PhcP.
36. A method as claimed in claim 5 wherein the DNA is positioned in frame with a constitutive promoter.
37. A method according to claim 36 wherein the constitutive promoter is Osmz or lacz.
38. A method as claimed in claim 3 or 4 wherein the DNA comprises DNA of SEQ ID No 7 or 8 or 9.
39. A method as claimed in claim 3 or 4 wherein the DNA comprises DNA of SEQ ID No 16.
40. A method as claimed in either of claims 3 or 4 wherein the vaccine comprises DNA of any one of the following SEQ ID Nos: 1, 3, 10.
41. A method as claimed in claim 4 wherein the DNA comprises DNA of SEQ ID No 20 or 22.
42. A method as claimed in claim 41 wherein the DNA is positioned down-stream of a eukaryotic promoter.
43. A method according to claim 42 wherein the eukaryotic promoter is a CMV immediate early promoter.
44. A method as claimed in claim 9 wherein the carrier is water.

45. A vaccine as claimed in claim 17 or 18 wherein the DNA comprises DNA of any of the sequences 7,8,9,10,16.
46. A vaccine as claimed in claims 19 or 45 wherein the DNA is positioned in frame with an in-vivo inducible promoter selected from one of the following: *htrA*, *nirB*, *ompR*, *ompC*, *phoP*.
47. A vaccine as claimed in claim 19 or 45 wherein the DNA is positioned in frame with a constitutive promoter selected from *Osmz* or *lacZ*.
48. A vaccine as claimed in claim 19 or 45 wherein the DNA is positioned downstream of a eukaryotic promoter.
49. A vaccine as claimed in claim 48 wherein the DNA comprises DNA of SEQ ID No 20 or 22.
50. A vaccine according to claim 23 wherein the carrier is water.

Fig.1.





Fig.2.



INTERNATIONAL SEARCH REPORT

Intern. Patent Application No.

PC 1/GB 96/00571

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/31 C07K14/24 C12N15/62 A61K39/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched: (classification system followed by classification symbols)

IPC 6 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	FEMS IMMUNOLOGY AND MEDICAL MICROBIOLOGY, 12 (3-4). 1995. 223-230., XP000573083 WILLIAMSON E D ET AL: "A new improved sub-unit vaccine for plague: The basis of protection" see the whole document	1-50
X	LEARY S E C ET AL: "Expression of Yersinia pestis V antigen in attenuated Salmonella typhimurium: Development of a novel vaccine for plague", KARGER AG, 13 (0). 1995. 216-217., BASEL, SWITZERLAND XP000572863 in Ravagnan G & Chiesa C (eds.): Yersiniosis: Present and Future. --- -/--	3,17,19

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier document but published on or after the international filing date
- "C" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search:

26 June 1996

Date of mailing of the international search report:

18.07.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 1
NL - 2380 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 631 epo nl,
Fax: (+31-70) 340-3016

Authorized officer:

Espin, J

INTERNATIONAL SEARCH REPORT

Inter- national Application No.
PCT/GS 96/00571

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>INFECTION AND IMMUNITY, vol. 63, no. 2, February 1995, WASHINGTON US, pages 563-568, XP002006749 OYSTON P C F ET AL.: "Immunization with live recombinant Salmonella typhimurium araA producing F1 antigen protects against plague" cited in the application see the whole document</p> <p>---</p>	3,7,19
X	<p>INFECTION AND IMMUNITY, vol. 62, no. 10, October 1994, WASHINGTON US, pages 4192-4201, XP002006750 MOTIN V L ET AL.: "Passive immunity to Yersinia mediated by anti-recombinant V antigen and protein A-V antigen fusion peptide" see the whole document</p> <p>-----</p>	3,17,19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB96/00571

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-14, 34-44
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 1-14, 34-44 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.